

The Antioxidant Effect of Noni Fruits Extract on HDL and LDL Level of Wistar Rat Induced by High Fat Diet

by Diah Purwaningsari, Sihning Ejt, Nita Pranitasari, Suhailah H., Lestari Dewi, Dian Ardiana

Submission date: 16-Feb-2023 07:32AM (UTC+0700)

Submission ID: 2015218216

File name: The_Antioxidant_Effect_of_Noni_Fruits_Extract_on_HDL_and.pdf (779.63K)

Word count: 3026

Character count: 16532

The Antioxidant Effect of Noni Fruits Extract on HDL and LDL Level of Wistar Rat Induced by High Fat Diet

Diah Purwaningsari¹, Sihning EJ Tehupuring², Nita Pranitasari¹, Suhailah Hayaza³, Lestari Dewi¹, Dian Ardiana¹

¹Faculty of Medicine, Hang Tuah University, Surabaya, Indonesia, ²Researcher, Faculty of Medicine, Hang Tuah University, Surabaya, Indonesia, ³Scientist and Data Analyst, Department of Biology, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia

Abstract

20

Dyslipidemia is an important risk factor for myocardial infarction, cardiovascular diseases, and Non-alcoholic Fatty Liver (NAFLD) disorders. Inflammatory conditions, oxidative stress, and insulin resistance, as well as several risk factors for cardiovascular disease, are also risk factors for this disease. This study aims to determine the effect of noni (*Morinda citrifolia*) extract on the improvement of dyslipidemia condition, based on the Low Density Lipoprotein (LDL) and High Density Lipoprotein (HDL) level. This research is a laboratory experimental study with a post-test only control group design. Male Wistar rat (*Rattus norvegicus*) were divided into four groups: negative control group with no induction and treatment (K-), positive control group given high-fat diet induction for 14 days (K+), noni extract treatment group given a high-fat diet and a noni extract at a dose of 100 mg BB per day for 7 days (P1), and noni extract at a dose of 200 mg/kg BB per day for seven days (P2). Low density Lipoprotein and High Density Lipoprotein serum levels were measured using the ELISA method. The results showed that there was a significant increase in LDL and a decrease in HDL in both treatment groups (P1 and P2) compared with K(+) group. Therefore, we conclude that noni extract (*Morinda citrifolia*) played a role as an antioxidant which can increase LDL and decrease HDL serum levels in rats induced by a high fat diet.

Keywords: Noni extract (*Morinda citrifolia*), high fat diet, LDL, HDL, antioxidant

Introduction

Dyslipidemia or atherogenic dyslipidemia is a term, which represents the occurrence of three things in humans, namely an increase in the concentration of Low Density Lipoprotein (LDL), a decrease in High Density Lipoprotein (HDL), and an increase in Triglycerides (TG). Dyslipidemia together with obesity, metabolic syndrome, insulin resistance, and type 2 diabetes mellitus are important risk factors for myocardial infarction and other cardiovascular diseases. Currently, several genes have been associated with this pattern of lipoprotein changes^[1]. Apart from cardiovascular disease, dyslipidemia can also lead to non-alcoholic

fatty liver (NAFLD) disease. Inflammatory conditions, oxidative stress and insulin resistance, as well as several risk factors for cardiovascular disease, are also risk factors for this disease^[2].

According to data, from 2009 to 2012, > 100 million US adults ≥20 years had a total cholesterol level of ≥200 mg / dL; nearly 31 million had levels ≥240 mg / dL. The data also show the presence of insulin resistance - which increases plasma triglyceride (TG) levels, an increase in Low Density Lipoprotein Cholesterol (LDL-C) and a decrease in High Density Lipoprotein Cholesterol (HDL-C) levels - which are important risk factors for vascular disease disorders^[3].

Riskesdas 2013 data shows, the prevalence in the population > 15 years of age, there is abnormal total cholesterol of 35.9 %, low HDL 22.9 %, LDL is not optimal with the combined category near optimal-

Corresponding author:

Diah Purwaningsari

email: diah.purwaningsari@hangtuah.ac.id

borderline high 60.3 % and high-very high category 15.9 %, abnormal triglycerides with high borderline category 13.0 % and high-very high category 11.9 %, and abnormal serum creatinine 6.0 %.

The currently established therapy of choice for dyslipidemia is statins. For conditions that are resistant to statins, a combination of statins with ezetimibe, fibrates, and nicotinic acid is used as a second choice. The next option is anti-PCSK9 drugs (evolocumab and alirocumab). In the latest development, to overcome homozygous FH (Familial Hypercholesterolemia), lomitapide and mipomersen therapy are used, but the price is still very expensive^[4].

Dyslipidemia requires long-term and repeated treatment, so the use of natural ingredients around us in the treatment of this disease is a very good breakthrough because drug side effects are expected to be minimal and costs incurred for treatment can be more efficient. Noni (*Morinda citrifolia*) or also called Noni, is a natural ingredient in the form of a plant with high antioxidant content, which is strong against free radicals, prevents oxidative damage to major biomolecules and provides significant protection against oxidative damage. This plant is rich in non-enzyme antioxidants such as ascorbic acid, α -tocopherol, carotenoids, flavonoids, phenols, tannins and carbohydrates. The contents of biological compounds in this plant include glycosides, polysaccharides, alkaloids, lignans, fattyacid esters, anthroquinones, scopoletin, morindin, vitamins and minerals, which can be extracted from the fruits, roots and leaves of Noni^[5].

Morinda citrifolia, or noni and locally known as noni, is a member of the madder family, Rubiaceae. This plant is native to tropical Asia and Australia and is distributed throughout the Pacific region. This plant is an evergreen shrub or small tree that grows to about 4–8 m. The branches are elongated short circular shape, thick, shiny ovoid leaves, dark green and muscular. The flowers, about 75–90 in number, are ovate with a round head. The length of the handle is 10–30 mm and the calyx is, in the form of a cut edge. White corolla, five lobes, with a greenish white tube, 7–9 mm^[6]. This fruit is used by people in all tropical regions of the world. The shiny green leaves are used by Caribs as a poultice for wounds, rheumatic joints, fevers and headaches. The

leaves are applied directly to the affected area for pain relief. Mashed fruit can also be applied directly to the affected area, including deep wounds and broken bones. It is also reported to cure various diseases ranging from arthritis, rheumatism, wounds, ulcers, and even to get rid of head lice^[6].

Antioxidants are substances that play an important role in inhibiting and scavenging free radicals, thus providing protection for humans against infections, degenerative diseases, and metabolic diseases. Much research has been focused on natural antioxidants derived from plants because they can produce safe therapies. Natural antioxidants can protect the human body from free radicals by slowing the progress of many chronic diseases as well as inhibiting dietary lipid oxidation. Many diseases related to oxidative stress are a result of the accumulation of free radicals in the body. Free radicals are involved in more than 80 diseases including Diabetes Mellitus, Arthritis, Cancer, Aging. Metabolic disease etc. In the treatment of this disease, antioxidant therapy is suspected to be the most important. In plants, there are free radical scavenger molecules that are rich in antioxidant activity. Free radicals are involved in more than 80 diseases including diabetes mellitus, arthritis, cancer, aging, and metabolic disease³. In the treatment of this disease, antioxidant therapy is suspected to be the most important. Therefore, this study was done to determine the effect of noni (*Morinda citrifolia*) extract on HDL and LDL serum levels in Wistar white rats (*Rattus norvegicus*) induced by a high-fat diet.

Materials and Methods

Experimental design

This study was a laboratory experimental research and used a post-test only control group design. There were four groups: two control groups and two treatment groups. Rats were randomly selected. This study was conducted in the Faculty of Medicine, Hang Tua University, Surabaya, Indonesia in 2019. The sample used in this study were 28 male white rats (*Rattus norvegicus*), aged 3–4 months with a bodyweight of \pm 150–200 grams, obtained from the Biochemistry Laboratory of the Medical Faculty of Hang Tuah University, Surabaya. The white rats were acclimatized for two weeks. They were maintained under a controlled temperature of $25\pm 2^\circ\text{C}$, humidity of $50\pm 10\%$, and a

12 h light-dark cycle and had free access to food and water. Animals were randomly selected and divided into 4 groups consisting negative control group with no induction and treatment (K-), a positive control group was given high-fat diet induction for 14 days (K+), noni extract treatment group given a high-fat diet and a noni extract at a dose of 100 mg/kg BB per day for 7 days (P1), and noni extract treatment group was given a high-fat diet and noni extract at a dose of 200 mg/kg BB per day for 7 days (P2).

Preparation of noni fruits crude extract

The noni fruits (*Morinda citrifolia*) were peeled off the outer skin, then split into several parts. After being chopped, the fruits were dried out using an oven at 60°C. The dried fruits were then mashed until it became powder. The powder was macerated with 70% ethanol for 24 hours and the filtrate was taken. Then, the filtrate was evaporated on the water heater^[7]. Noni extract was obtained.

Preparation of Pulvis Gummi Arabicum solution

The 3% PGA solution was used in the negative control group during the treatment stage. Hot water was poured into the mortar. Around three grams of PGA were sprinkled and inflated for 15 minutes. It was stirred until homogeneous. Then, 100 mL of distilled water was added.

Blood serum analysis

On the 29th day, the rat's blood was collected. The rats were previously was fasten for ± 12 hours and were only given water to drink. Then the rats were anesthetized and blood was drawn from the rats' hearts (intracardiac) using a syringe. Each rat had approximately 3 ml of blood. Then the blood was collected in a serological container without anticoagulants and closed with a rubber cap. The sample proceeded for HDL and LDL levels analysis.

HDL and LDL level analysis

Blood samples were collected in tubes and then centrifuged at 3000× g for 20 min. The serum was

obtained. The levels of HDL and LDL in the serum were performed using an ELISA kit (Bioassay Technology, Shanghai, China) according to the manufacturer's protocol. After adding 40µl blood serum to wells, around 10µl of anti-HDL and anti-LDL, and 50µl streptavidin-HRP was added. Plates were incubated for 60 minutes at 37°C and washed 5 times with wash buffer. Then, 50µl substrate solution A and B were added to each well and incubated for 10 minutes at 37°C in the dark. Lastly, a 50µl stop solution was added to each well. The optical density was measured using a microplate reader ($\lambda=450$ nm).

Statistical Analysis

Data from experiments were expressed as mean \pm standard deviation (SD). Statistical analyses were performed by one-way analysis of variance (ANOVA) followed by Duncan's post hoc test to compare the groups using SPSS version 21 (SPSS Inc., Chicago, IL). The significance level for analyses was set at $p < 0.05$.

Result and Discussion

Noni extract contains several active ingredients, namely phenolic compounds and flavonoid, which function as antioxidants and reduce levels of lipid oxidation and increase catalase activity. The main flavonoids that have been detected in *Morinda citrifolia* are catechins and epicatechins^[8]. Inhibition of active ingredients of phenolic and flavonoid compounds will cause inhibition of Reactive Oxygen Species, which are produced from modified lipoprotein protein molecules due to oxidation, glycosylation and glycosidation processes^[9].

In this study, 5% induction of a high-fat diet for 14 days was proven to increase the LDL blood serum levels of male Wistar white rats (*Rattus norvegicus*), and administration of noni extract (*Morinda citrifolia*) at a dose of 100 mg/kg BW for 7 days was proven to be effective in lowering LDL levels (Table 1). Rats given a high-fat diet will experience an increase in LDL levels, this is due to the intake of saturated fatty acids contained in goat fat, lard, egg yolks and cholic acid^[10].

Table 1. Effect of noni fruits on HDL and LDL level in rats induced by a high-fat diet. *p<0.05 compared to the positive control group. **p<0,05 compared to both control groups.

Groups	LDL level (µg/mL)	HDL Level (µg/mL)
Negative control	6,86 ± 1,86	31 ± 2,16
Positive control	10,72 ± 1,38	25,86 ± 1,22
P1	5,86 ± 1,35*	30,29 ± 7,72*
P2	5,29 ± 0,95*	33,29 ± 7,50*

Other studies that are in line with this study are research conducted in vitro on animals and active smokers, and the protective effect of noni (*Morinda citrifolia*) / Noni Fruit juice on superoxide anion radicals and lipid peroxide. The effect of this noni fruit can contribute to lowering lipid levels. A double-blind placebo controlled study conducted in a small population showed a decrease in total cholesterol and triglyceride levels in Noni Fruit drinkers^[11].

The study stated that the treatment group, namely the group given Noni fruit juice at a dose of 50 and 100 mg/kg BW showed a decrease in total cholesterol, triglycerides and VLDL-cholesterol at both doses when compared to control the disease. However, the reduction in total cholesterol and LDL-C in the administration of noni fruit juice at a dose of 50 mg/kg BW failed to show a statistically significant reduction compared to atorvastatin, a cholesterol-lowering drug that has been established. However, this study has sufficiently proven the hypolipidemic activity of Noni fruit juice in a high-fat diet induced hyperlipidemia in rats^[11].

The study, which was conducted on 20 mice, used 40% sucrose induction for 10 weeks to make a metabolic drome model, then gave *Morinda citrophilia* extract at a dose of 200 mg/kg BW for 2 weeks in the treatment group compared to the standard diet. Observations were made on rat body weight, serum glucose levels, total cholesterol levels, HDL, TG, uric acid, total protein (PTOT), albumin and globulin. The results showed that the consumption of water extract from dried *M. citrifolia* leaves reduced abdominal fat and triacylglycerols which are associated with increased adiponectin levels and

changes in adipose tissue cell dynamics^[12].

Many diseases related to oxidative stress are a result of the accumulation of free radicals in the body^[13]. The main mechanism of antioxidants maintained in mammals is through the neutralization of ROS with glutathione (GSH) and three main scavenging antioxidant enzymes, namely superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GSH-Px). Besides, matrix metalloproteinases (MMPs) promote the degradation of the extracellular matrix (ECM) molecules that play a role in the development of liver damage or even fibrosis^[14,15].

Morinda citrifolia can prevent free radical-induced oxidative-pathological events in the liver by inhibiting the inflammatory response and suppressing increased liver enzyme activity; thus preventing damage due to cell membranes. The addition of *Morinda citrifolia* extract shows hepatoprotection against alcohol-induced injury due to regulation of lipid homeostasis, antioxidant status, and alcohol metabolism^[15].

Conclusion

In sum, the administration of noni fruit extracts have successfully increased LDL serum level and lowered HDL serum levels in dyslipidemia Wistar white rats induced by a high-fat diet.

Conflict of Interest : The authors declare that they have no conflict of interest.

Source of Funding : This study was supported by Hang Tuah University, Surabaya, Indonesia.

Acknowledgements : Authors would like to thank

Hang Tua University for giving support and funding toward this research.

23

Ethical Approval: Ethical approval for this study was obtained from the Committee of Animal Care

28

Use, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia (approval No. 541-KE).

References

- Musunuru K. Atherogenic dyslipidemia: cardiovascular risk and dietary intervention. *Lipids*. 2010; 45: 907-914.
- Katsiki N, Mikhailidis DP, Mantzoros CS. Non-alcoholic fatty liver disease and dyslipidemia: An update. *Metabolism: clinical and experimental*. 2016; 65: 1109-1123.
- Jellinger PS, Handelsman Y, Rosenblit PD, et al. American Association of Clinical Endocrinologists and American College of Endocrinology Guidelines for Management of Dyslipidemia and Prevention of Cardiovascular Disease. *Endocrine practice : official journal of the American College of Endocrinology and the American Association of Clinical Endocrinologists*. 2017; 23: 1-87.
- Zodda D, Giammona R, Schifilliti S. Treatment Strategy for Dyslipidemia in Cardiovascular Disease Prevention: Focus on Old and New Drugs. *Pharmacy*. 2018; 6(1): 10.
- Durairaj VSaB. Antioxidant And Free Radical Scavenging Effect Of Morinda Citrifolia Fruit Extract. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2014; 6: 55-59.
- Rosalizan M, Rohani M, Khatijah I. Physico-chemical characteristics of *Morinda citrifolia* fruit during growth and maturation. *J. Trop. Agric. Food Sci*. 2010; 38: 21-30.
- Hayaza S, Istiqomah S, Susilo RJK, et al. Antidiabetic activity of ketapang (*Terminalia catappa* L.) leaves extract in streptozotocin-induced diabetic mice. *Indian Vet J*. 2019; 96: 11-13.
- Mohd Zin Z, Abdul Hamid A, Osman A, et al. Isolation and Identification of Antioxidative Compound from Fruit of Mengkudu (*Morinda citrifolia* L.). *International Journal of Food Properties*. 2007; 10(2): 363-373
- Gupta RK, Singh N. *Morinda citrifolia* (noni) alters oxidative stress marker and antioxidant activity in cervical cancer cell lines. *Asian Pacific Journal of Cancer Prevention*. 2013;14: 4603-4606.
- Adiputro DL, Widodo MA, Romdoni R, et al. Extract of mangosteen increases high density lipoprotein levels in rats fed high lipid. *Universa Medicina*. 2015; 32: 37-43.
- Shoeb A, Alwar MC, Shenoy PJ, et al. Effect of *Morinda citrifolia* (Noni) Fruit Juice on High Fat Diet Induced Dyslipidemia in Rats. *Journal of clinical and diagnostic research : JCDR*. 2016; 10: FF06-10.
- Ida SR, Rodolfo QC, Jorge RA, et al. Beneficial effects of *Morinda citrifolia* Linn.(noni) leaf extract on obesity, dyslipidemia and adiponectinemia in rats with metabolic syndrome. *International Journal of Pharmaceutical Sciences and Research*. 2017; 8: 2496-2503.
- Purnamasari R, Winarni D, Permanasari AA, et al. Anticancer activity of methanol extract of *Ficus carica* leaves and fruits against proliferation, apoptosis, and necrosis in Huh7it cells. *Cancer Informatics*. 2019; 18: 1-7.
- Purwaningsari D, Nugraha J, Wahyuningsih SPA, et al. Effect of Polysaccharide Krestin on MMP3 Expression and Foot Diameter in Rheumatoid Arthritis in Rat. *Indian Vet. J*. 2020; 97(01): 24-26.
- Soto-Rodríguez I, Quintana-Castro R, Cano-Martínez CA, Aguirre-Maldonado I, Granados-Echegoyen C, Alexander-Aguilera A. *Morinda citrifolia* (noni) changes the expression of AdipoR2 mRNA in hepatic tissue of rats with nonalcoholic fatty liver disease (NAFLD). *Journal of Pharmacognosy and Phytochemistry*. 2017; 6: 120-125.

The Antioxidant Effect of Noni Fruits Extract on HDL and LDL Level of Wistar Rat Induced by High Fat Diet

ORIGINALITY REPORT

10%

SIMILARITY INDEX

5%

INTERNET SOURCES

8%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 1 | "Association between Dietary Pattern and Incidence of Cholesterolemia in Korean Adults: The Korean Genome and Epidemiology Study", <i>Nutrients</i> , 2018
Publication | <1 % |
| 2 | Submitted to Barry University
Student Paper | <1 % |
| 3 | Hyeonkang Park, Chang-Soo Han, Chun-Woong Park, Kibum Kim. "Change in spray behavior of pharmaceutical particles by creating bypass hole in dry powder inhaler", <i>Powder Technology</i> , 2022
Publication | <1 % |
| 4 | Normah Ismail, Nurulain Abd Razak. "Characterization and Purification of Protease Extracted from Two Maturity Stages of 'Noni' (<i>Morinda citrifolia</i> L.) fruit", <i>Scientific Research Journal</i> , 2014
Publication | <1 % |

5

Chang Shan, Deng Zhang, Dong-ni Ma, Yan-fang Hou et al. "Osteocalcin ameliorates cognitive dysfunctions in a mouse model of Alzheimer's Disease by reducing amyloid β burden and upregulating glycolysis in neuroglia", Cell Death Discovery, 2023

Publication

<1 %

6

acervodigital.unesp.br

Internet Source

<1 %

7

translational-medicine.biomedcentral.com

Internet Source

<1 %

8

www.jove.com

Internet Source

<1 %

9

Yang Li, Wen-Jun Ma, Bao-Kun Qi, Sami Rokayya, Dan Li, Jing Wang, Hong-Xia Feng, Xiao-Nan Sui, Lian-Zhou Jiang. "Blending of Soybean Oil with Selected Vegetable Oils: Impact on Oxidative Stability and Radical Scavenging Activity", Asian Pacific Journal of Cancer Prevention, 2014

Publication

<1 %

10

Joon-Kwan Moon, Takayuki Shibamoto. "Antioxidant Assays for Plant and Food Components", Journal of Agricultural and Food Chemistry, 2009

Publication

<1 %

- | | | |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 11 | Meilisa Tabrani, Djunizar Djamaludin.
"Pemberian Jus Apel Hijau Terhadap
Penurunan Kadar Kolesterol Pada Pasien
Hiperkolesterolemia Di Kota Metro", JURNAL
KREATIVITAS PENGABDIAN KEPADA
MASYARAKAT (PKM), 2021
Publication | <1 % |
| 12 | csj.cumhuriyet.edu.tr
Internet Source | <1 % |
| 13 | discovery.ucl.ac.uk
Internet Source | <1 % |
| 14 | medplants.blogspot.com
Internet Source | <1 % |
| 15 | myresearchspace.uws.ac.uk
Internet Source | <1 % |
| 16 | www.intechopen.com
Internet Source | <1 % |
| 17 | www.pubfacts.com
Internet Source | <1 % |
| 18 | Jie Jing, Shiling Chen, Xuan Wu, Jingfei Yang,
Xia Liu, Jiahui Wang, Jingyi Wang, Yunjie Li,
Ping Zhang, Zhouping Tang. "rtPA Directly
Protects Neurons After Intracerebral
Hemorrhage through PI3K/AKT/mTOR
Pathway", Cold Spring Harbor Laboratory,
2023 | <1 % |

19

Lili Liu, Hong Li, Yang Zhang, Jing Zhang, Zhenhuan Cao. "Hepatitis B virus infection combined with nonalcoholic fatty liver disease: Interaction and prognosis", Heliyon, 2023

Publication

<1 %

20

Rodolfo Quintana Castro, Isaac Aguirre Maldonado, Ida Soto Rodríguez, Rosa A. Deschamps Lago et al. "Cd36 gene expression in adipose and hepatic tissue mediates the lipids accumulation in liver of obese rats with sucrose-induced hepatic steatosis", Prostaglandins & Other Lipid Mediators, 2019

Publication

<1 %

21

www.krishisanskriti.org

Internet Source

<1 %

22

Alkilany Salem Abuzaid, Elin Yulinah Sukandar, Neng Fisher Kurniati, I. Ketut Adnyana. "Antihyperlidemic Effects of Mangosteen (Garcinia mangostana L.) Pericarp Ethanolic Extract In High-Carbohydrate Wistar Rats", JOURNAL OF NATURAL REMEDIES, 2017

Publication

<1 %

23

Naohito HIBINO, Masahiro YAMANO, Yoshitaka HAMADA, Yoshinori TAKAHASHI. "Use of a Free Medial Femoral Condyle Periosteal Flap in Digital Reconstruction – A

<1 %

Report of Two Patients", The Journal of Hand Surgery (Asian-Pacific Volume), 2022

Publication

24

Rakesh Kumar Gupta, Neeta Singh. "Morinda citrifolia (Noni) Alters Oxidative Stress Marker and Antioxidant Activity in Cervical Cancer Cell Lines", Asian Pacific Journal of Cancer Prevention, 2013

Publication

<1 %

25

Amy C. Brown. "Anticancer Activity of Morinda citrifolia (Noni) Fruit: A Review", Phytotherapy Research, 2012

Publication

<1 %

26

Buthaina Aamir Ali Al-Rahbi, Abdullah Mohammed Al-Sadi, Majida Mohammed Ali Al-Harrasi, Jamal Nasser Al-Sabahi et al. "Effectiveness of Endophytic and Rhizospheric Bacteria from Moringa spp. in Controlling Pythiumaphanidermatum Damping-Off of Cabbage", Plants, 2023

Publication

<1 %

27

Tingting Xin, Jun Li, Qian Xu. "Histological and bone mineral density analyses of the mandibular lamellar/oblique fracture healing in beagles.", Research Square Platform LLC, 2023

Publication

<1 %

28

notablesdelaciencia.conicet.gov.ar

Exclude quotes On
Exclude bibliography On

Exclude assignment template On
Exclude matches Off