

The Effect of Brown Seaweed (Sargassum sp.) Extract on Apoptosis Process in Breast Cancer – A Literature Review

by Ismylatifa Devi, Diah Purwaningsari, Nita Pranitasari, Herin
Setianingsih,

Submission date: 16-Feb-2023 07:33AM (UTC+0700)

Submission ID: 2015218876

File name: The_Effect_of_Brown_Seaweed_Sargassum_sp._Extract_on.pdf (794.58K)

Word count: 2729

Character count: 15615

The Effect of Brown Seaweed (*Sargassum sp.*) Extract on Apoptosis Process in Breast Cancer – A Literature Review

Ismylatifa Devi¹, Diah Purwaningsari², Nita Pranitasari², HerinSetianingsih²

¹Student at Faculty of Medicine, Hang Tuah University, Surabaya, Indonesia, ²Lecturers at Faculty of Medicine, Hang Tuah University, Surabaya, Indonesia

Abstract

Breast cancer was one of the most common types of cancer in women and the major cause of death. Chemotherapy in breast cancer caused toxicity to the normal cells, resistance to the drugs, and several other side effects. Seaweed had many functional compounds that could be used as adjuvant therapy in breast cancer because of its anti-cancer properties called the proapoptotic agent. These compounds were fucoidan, phloroglucinol, and fucoxanthin. Their proapoptotic mechanisms shown by the journal were through increasing proapoptotic protein expression, decreasing antiapoptotic protein expression, inducing oxidative stress production on the cancer cell, and inhibiting the PI3K pathway. This review concludes that the brown seaweed (*Sargassum sp.*), which contains fucoidan, phloroglucinol, and fucoxanthin, could increase breast cancer cells' apoptotic process.

Keywords: Apoptosis, Breast Cancer, Brown Seaweed, Oxidative Stress, Proapoptotic, *Sargassum sp.*

Introduction

Breast cancer is the most prevalent cancer in women, there were 2.1 million women every year suffered breast cancer, and in 2018 around 627,000 women died from breast cancer⁽¹⁾. Nowadays, breast cancer's main treatments are surgery, chemotherapy, radiation, hormone therapy, and targeted therapy. These treatments are beneficial, although they can cause serious side effects such as nausea, bone marrow failure, resistance to drugs, and another significant clinical toxic effects^(2,3). Recent cancer studies have focused on designing drugs from natural sources because of the fewer side effects than synthetic anti-cancer drugs. A new strategy has been proposed to reduce the dosage of synthetic drugs by providing a combination of natural compounds; one of them was seaweed. In China and Japan, several traditional medicines utilize seaweed to treat tumors^(4,5).

Sargassum is a brown seaweed genus with a tropical and subtropical distribution, which exists in all oceans. *Sargassum sp.* is a type of seaweed that is spread in Indonesia^(6,7). Brown seaweed contains bioactive compounds such as fucoidan, phloroglucinol and fucoxanthin. Several studies show they have anti-cancer activity, through inducing apoptosis in cancer cells⁽⁴⁾.

Apoptosis is a natural mechanism of cell death, and there are two mechanisms, the intrinsic pathway using mitochondrial proteins and the extrinsic pathway using extracellular signal induction. Apoptosis is very influential on cancer, so it is used as one of cancer treatment targets. Reactive Oxygen Species (ROS) or free radicals also play an essential role in the initiation, promotion and development of cancer, so antioxidants are needed to prevent cancer. But recent studies are interested in pro-oxidants as an anti-cancer therapy by inducing apoptosis with mitochondrial-dependent pathways and involvement of ROS induction⁽⁸⁾. This research was conducted to determine the effect of brown seaweed (*Sargassum sp.*) on the apoptotic process of breast cancer cells.

Corresponding Author :

Diah Purwaningsari

email: diah.purwaningsari@hangtuah.ac.id

Methodology

Articles were collected using Pubmed and Science Direct database. The search words include apoptosis, breast cancer, brown seaweed, *Sargassum sp.*, oxidative stress, PI3K pathway, and proapoptotic. Articles were collected from the year 2015-2020 and indexed in Scimago and Scopus.

Effect of fucoidan, phloroglucinol, and fucoxanthin on apoptosis

³ Fucoidan is a fucose-enriched and sulfated polysaccharide obtained from the extracellular matrix of brown seaweed. Fucoidan has several anti-cancer mechanisms; one of them is inducing apoptosis⁽⁹⁾. Table 1 show some research about fucoidan anti-cancer activities. Fucoidan extract of *F. vesiculosus* was tested on different female cancer cell lines (breast, ovarian, uterine, endometrial carcinoma) in vitro. In most cancer cells, fucoidan treatment decreased in phosphorylated PI3K, AKT and mTOR. Fucoidan (*U. pinnatifida*) inhibits PI3K / AKT phosphorylation in prostate cancer cells in vitro. After treatment with fucoidan, the AML (Acute Myeloid Leukemia) cell line decreased phosphorylated AKT by the in vitro method. The same decrease in AKT was seen in the other two AML cell lines (NB4 and HL60) when treated with fucoidan in vitro, derived from *F. vesiculosus*⁽¹⁰⁾.

Phloroglucinol, one of the derivatives of phlorotannin has no toxicity but has a more protective role in normal

tissue cells. It was found in a study that phloroglucinol showed anti-cancer properties through upregulation of p53, Bax activation, Bcl-2 inhibition, increased caspase 3 and 9 activity and was associated with downregulation of the NF- κ B line. Phloroglucinol can inhibit the growth of MCF-7 cells which induces apoptosis through the mitochondrial pathway with reactive oxygen species (ROS) production⁽⁴⁾.

Fucoxanthin is major carotenoids produced by seaweeds, and their structure includes an allelic bond and an oxygenic functional group; these compounds have both protective and photosynthetic functions⁽¹¹⁾. Several studies have reported that fucoxanthin's anti-cancer properties and its metabolites are pro-oxidants that can trigger apoptosis. A study has observed growth inhibition in the leukaemia cell lines by fucoxanthin and has linked it to ROS formation by fucoxanthin leading to apoptosis. There was an increase in H₂O₂ and O₂⁻ production due to treatment with fucoxanthin and the accumulation of cells containing sub-G₁ DNA content (indicating cessation of the cell cycle in stage G₁). In co-treatment with the commercial antioxidant NAC (N-acetylcysteine), the number of apoptotic bodies and cell DNA fragmentation decreased, which correlated the apoptotic effect fucoxanthin with the resulting ROS. Thus, they concluded that fucoxanthin's cytotoxic effect through ROS formation triggered apoptosis in HL-60 cells⁽¹²⁾. Several studies show fucoxanthin's anticancer activities in table 1.

Table 1. Effect of fucoidan, phloroglucinol, and fucoxanthin on apoptosis

Bioactive Compound	Method	Dosage	Sample	Result	Mechanism	Ref.
Fucoidan	In vitro	6.25, 12.5, and 25 μ g/mL for 48 hours	MDA-MB-231 cell	Increase apoptosis process	Inhibit PI3K pathway	(13)
	In vivo	200 and 400 mg/kg.bw for 16 weeks	Sprague Dawley mice induced breast cancer			
	In vitro	0.8mg/mL for 48 hours	Melanoma B16 cell		Regulate apoptosis protein expression and increase ROS production	(14)
	In vivo	200 and 400 mg/kg.bw for 3 days	Sprague Dawley mice injected MCF-7 breast cancer cells		Modulate E-cadherin expression	(3)
	In vitro	50 and 150 μ g/mL for 24 hours	MCF-7 cell		Activate caspase-8	(15)
	In vivo	500 and 1000 mg/kg.bw for 24 hours	HCC SMMC-7721 cell		Increase ROS production and activate caspase 3 and 9	(5)

Table 1. Effect of fucoidan, phloroglucinol, and fucoxanthin on apoptosis

Phloroglucinol	In vitro	10 and 30 μ M for 48 hours	MCF-7 cell	Increase apoptosis process	Regulate apoptosis protein expression and activate caspase 3 and 9	(16)
	In vivo	25mg/kg.bw for 3 days	BALB / C mice injected breast cancer cells BT549 and MDA-MB231		Inhibit RAS signaling pathway	(4)
Fucoxanthin	In vitro	10 μ M for 24 hours	MDA-MB-231 and MCF-7 cell	Increase apoptosis process	Regulate apoptosis protein expression and activate caspase pathway	(17)
	In vivo	200mg/kg.bw for 28 days	BALB / C mice injected Glioma U87 and U251 cell		Regulate apoptosis protein expression and inhibit PI3K/ AKT/m-TOR and p38	(18)
	In vitro	25 and 50 μ M for 24 hours	Glioma U87 and U251 cell		Regulate apoptosis protein expression	

Role of oxidative stress on apoptosis

Oxidative stress refers to an imbalance between prooxidant and antioxidant factors controlled by many components; this imbalance can cause cell damage. ROS plays a vital role in oxidative stress and is produced as a byproduct of cellular metabolism, particularly in mitochondria. Once accumulated, ROS can attack cellular proteins, DNA, and lipids, leading to a state of oxidative stress. ROS contributes to several human diseases, including cardiovascular, inflammatory and neurodegenerative diseases, and cancer. Increased levels of mitochondrial ROS are sufficient to induce apoptosis⁽⁸⁾.

The polysaccharide extract of *Sargassum wightii* was given to MCF-7 and MDA-MB-231 breast cancer cells. Polysaccharides significantly reduce the growth of both breast cancer cells, depending on the dose. The inhibition of polysaccharide growth against MCF7 cells was 69% at 500 μ g/ml concentration. Furthermore, the inhibition rate of polysaccharide growth against MDA-MB-231 cells was 73% at 500 μ g/ml concentration. The results showed polysaccharides induced apoptosis in breast cancer cells by increasing ROS formation, mitochondrial membrane division and nuclear destruction. Polysaccharides also increase caspase 3/9 activity, causing apoptosis in breast cancer cells. Based on this study, polysaccharides from *Sargassum wightii* can be candidates for further evaluation as anti-cancer agents for cancer in humans, particularly breast cancer⁽¹⁹⁾.

Previous studies have reported the potential of polysaccharides from *Sargassum fusiforme* to induce apoptosis in leukaemia, stomach, bladder, and breast cancer cells. This study aims to determine *S. fusiforme* polysaccharides' effect on apoptosis and regulation of Human Erythroleukaemia (HEL) cells. The effect on HEL cell growth was detected by the Cell Counting Kit-8 (CCK-8) method, and apoptosis was detected by Hoechst staining. Cell cycle distribution and apoptosis were identified using flow cytometry⁽⁸⁾. Cell cycle gene expression, p53, antiapoptosis gene, Bcl-xL and Bcl-2, and pro-apoptosis, Bax, Bad, and caspase-3 genes, as well as the corresponding protein expression, were detected using qPCR (quantitative Polymerase Chain Reaction) and western blot. The results showed that it decreased the viability of HEL cells and induced apoptosis of HEL cells. The apoptosis induction mechanism was found to be related to the formation of ROS, which in turn can lead to DNA damage and cell death, associated with induction of cell cycle arrest in the G₀ / G₁ phase, and increased expression of apoptosis-related genes and proteins⁽²⁰⁾.

Role of PI3K pathway on apoptosis

The PI3K signaling pathway is the center of the intracellular signaling axis that integrates multiple signals to induce cancer cells' growth and development. In general, the signaling path includes three main components consisting of PI3K, AKT and m-TOR. PI3K

is a kinase enzyme family that catalyzes PIP2 to form PIP3. PIP3 production promotes AKT phosphorylation as the primary regulator that can directly modulate several downstream targets, including the m-TOR and WNT / β -catenin pathways. AKT can regulate cancer cell proliferation, metabolism, apoptosis and angiogenesis. The m-TOR complex is activated by AKT and regulates several cell growth functions, including protein synthesis, cell survival, and autophagy inhibition. Recent data show that PI3K signaling is frequently activated in several malignancies in humans, including breast cancer and prolonged activation of PI3K signaling is associated with poor prognosis and resistance to chemotherapy⁽²¹⁾.

Mutations in this signaling pathway are common in cancers, particularly in breast cancer. About 60% of tumours have genetic changes that cause hyperactivation of the PI3K / AKT / m-TOR pathway. Preclinical data from cell-based and genetic engineering studies (GEMs) have demonstrated that this mutation activates PI3K / AKT / m-TOR signaling and is an oncogenic driver by promoting cell transformation, tumour initiation, development, and resistance to apoptosis⁽²²⁾. Cancer cells have developed strategies to avoid apoptosis, mutations in p53 and decreased expression of a molecule called caspase which is the crucial mediator and effector of apoptosis, disrupting external signaling and altering the balance between pro and anti-apoptotic molecules. Clinical studies in cancer have shown that targeting apoptosis is an effective strategy in fighting cancer. One strategy in targeting apoptosis is to target peripheral pathways effective in cell survival and apoptosis, such as PI3K-Akt-mTOR⁽²³⁾. Several studies in table 1 support this strategy.

Conclusion

Brown seaweed (*Sargassum sp.*) which contains fucoidan, phloroglucinol and fucoxanthin can increase the apoptosis process in breast cancer with several mechanisms, through regulating apoptotic protein, increasing proapoptotic protein and decreasing anti-apoptotic protein, inducing oxidative stress (ROS) production and inhibiting the PI3K pathway.

Ethical Clearance – Not required since it is a literature review

Source of Funding – Nil

Conflict of Interest – Nil

References

1. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2018;68:398.
2. Palanisamy S, Vinosha M, Manikandakrishnan M, Anjali R, Rajasekar P, Marudhupandi T, et al. Investigation of antioxidant and anticancer potential of fucoidan from *Sargassum polycystum*. *Int J Biol Macromol.* 2018;116:151–61.
3. He X, Xue M, Jiang S, Li W, Yu J, Xiang S. Fucoidan promotes apoptosis and inhibits EMT of breast cancer cells. *Biol Pharm Bull.* 2019;42(3):442–7.
4. Pádua D, Rocha E, Gargiulo D, Ramos AA. Bioactive compounds from brown seaweeds: Phloroglucinol, fucoxanthin and fucoidan as promising therapeutic agents against breast cancer. *Phytochem Lett.* 2015;14:91–8.
5. Sanjeewa KKA, Lee JS, Kim WS, Jeon YJ. The potential of brown-algae polysaccharides for the development of anticancer agents: An update on anticancer effects reported for fucoidan and laminaran. *Carbohydr Polym.* 2017;3–11.
6. Máximo P, Ferreira LM, Branco P, Lima P, Lourenço A. Secondary metabolites and biological activity of invasive macroalgae of southern Europe. *Mar Drugs.* 2018;16(8):1–28.
7. Puspita M, Setyawidati NAR, Stiger-Pouvreau V, Vandanon L, Widowati I, Radjasa OK, et al. Indonesian *Sargassum* species bioprospecting: potential applications of bioactive compounds and challenge for sustainable development. *Adv Bot Res.* 2020;95:113–61.
8. León-González AJ, Auger C, Schini-Kerth VB. Pro-oxidant activity of polyphenols and its implication on cancer chemoprevention and chemotherapy. *Biochem Pharmacol.* 2015;98(3):371–80.
9. Wu J, Li H, Wang X, Zhang X, Liu W, Wang Y, et al. Effect of polysaccharide from *Undaria pinnatifida* on proliferation, migration and apoptosis of breast cancer cell MCF7. *Int J Biol Macromol.* 2019;121:734–42.
10. van Weelden G, Bobi M, Okla K, van Weelden WJ,

- Romano A, Pijnenborg JMA. Fucoidan structure and activity in relation to anti-cancer mechanisms. *Mar Drugs*. 2019;17(32):1–30.
11. Gutiérrez-Rodríguez AG, Juárez-Portilla C, Olivares-Bañuelos T, Zepeda RC. Anticancer activity of seaweeds. *Drug Discov Today [Internet]*. 2018;23(2):434–47.
12. Saini RK, Keum YS, Daglia M, Rengasamy KR. Dietary carotenoids in cancer chemoprevention and chemotherapy: A review of emerging evidence. *Pharmacol Res*. 2020;157:1–12.
13. Xue M, Ji X, Xue C, Liang H, Ge Y, He X, et al. Caspase-dependent and caspase-independent induction of apoptosis in breast cancer by fucoidan via the PI3K/AKT/GSK3 β pathway in vivo and in vitro. *Biomed Pharmacother*. 2017;94:898–908.
14. Sanjeewa KKA, Kang N, Ahn G, Jee Y, Kim YT, Jeon YJ. Bioactive potentials of sulfated polysaccharides isolated from brown seaweed *Sargassum* spp in related to human health applications: A review. *Food Hydrocoll*. 2018;81:200–8.
15. Palanisamy S, Vinosha M, Marudhupandi T, Rajasekar P, Prabhu NM. Isolation of fucoidan from *Sargassum polycystum* brown algae: Structural characterization, in vitro antioxidant and anticancer activity. *Int J Biol Macromol*. 2017;102:405–12.
16. Kim RK, Suh Y, Yoo KC, Cui YH, Hwang E, Kim HJ, et al. Phloroglucinol suppresses metastatic ability of breast cancer cells by inhibition of epithelial-mesenchymal cell transition. *Cancer Sci*. 2015;1–8.
17. Rwigemera A, Mamelona J, Martin LJ. Comparative effects between fucoxanthinol and its precursor fucoxanthin on viability and apoptosis of breast cancer cell lines MCF-7 and MDA-MB-231. *Anticancer Res*. 2015;35:207–20.
18. Liu Y, Zheng J, Zhang Y, Wang Z, Yang Y, Bai M, et al. Fucoxanthin Activates Apoptosis via Inhibition of PI3K/Akt/mTOR Pathway and Suppresses Invasion and Migration by Restriction of p38-MMP-2/9 Pathway in Human Glioblastoma Cells. *Neurochem Res*. 2016;41(10):2728–51.
19. Vaikundamoorthy R, Krishnamoorthy V, Vilwanathan R, Rajendran R. Structural characterization and anticancer activity (MCF7 and MDA-MB-231) of polysaccharides fractionated from brown seaweed *Sargassum wightii*. *Int J Biol Macromol*. 2018;111:1229–37.
20. Ding HM, Chen XJ, Chen HM, Wang CS, Qian GY. Effect of *Sargassum fusiforme* polysaccharide on apoptosis and its possible mechanism in human erythroleukemia cells. *Chin J Nat Med*. 2020;18(10):749–59.
21. Rahmani F, Ferns GA, Talebian S, Nourbakhsh M, Avan A, Shahidsales S. Role of regulatory miRNAs of the PI3K/AKT signaling pathway in the pathogenesis of breast cancer. *Gene*. 2020;737:144459.
22. Guerrero-Zotano A, Mayer IA, Arteaga CL. PI3K/AKT/mTOR: role in breast cancer progression, drug resistance, and treatment. *Cancer Metastasis Rev*. 2016;35(4):515–24.
23. Mirza-Aghazadeh-Attari M, Ekrami EM, Aghdas SAM, Mihanfar A, Hallaj S, Yousefi B, et al. Targeting PI3K/Akt/mTOR signaling pathway by polyphenols: Implication for cancer therapy. *Life Sci*. 2020;255(2020):1–21.

The Effect of Brown Seaweed (Sargassum sp.) Extract on Apoptosis Process in Breast Cancer – A Literature Review

ORIGINALITY REPORT

7 %

SIMILARITY INDEX

4 %

INTERNET SOURCES

5 %

PUBLICATIONS

1 %

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|--|
| <div style="background-color: red; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 0 auto;">1</div> | <p>Subramanian Palanisamy, Manoharan Vinosha, Muthushanmugam Manikandakrishnan, Ravichandran Anjali et al. "Investigation of antioxidant and anticancer potential of fucoidan from Sargassum polycystum", International Journal of Biological Macromolecules, 2018</p> <p>Publication</p> | <div style="font-size: 2em;">1</div> % |
| <hr/> | | |
| <div style="background-color: magenta; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 0 auto;">2</div> | <p>Angel Guerrero-Zotano, Ingrid A. Mayer, Carlos L. Arteaga. "PI3K/AKT/mTOR: role in breast cancer progression, drug resistance, and treatment", Cancer and Metastasis Reviews, 2016</p> <p>Publication</p> | <div style="font-size: 2em;">1</div> % |
| <hr/> | | |
| <div style="background-color: purple; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 0 auto;">3</div> | <p>Lei Wu, Jing Sun, Xitong Su, Qiuli Yu, Qiuyang Yu, Peng Zhang. "A review about the development of fucoidan in antitumor activity: Progress and challenges", Carbohydrate Polymers, 2016</p> <p>Publication</p> | <div style="font-size: 2em;">1</div> % |
-

4	pericles.pericles-prod.literatumonline.com Internet Source	1 %
5	www.transparencymarketresearch.com Internet Source	1 %
6	Kalu Kapuge Asanka Sanjeewa, Nalae Kang, Ginnae Ahn, Youngheun Jee, Young-Tae Kim, You-Jin Jeon. "Bioactive potentials of sulfated polysaccharides isolated from brown seaweed Sargassum spp in related to human health applications: A review", Food Hydrocolloids, 2018 Publication	1 %
7	www.genox.com Internet Source	1 %
8	mobt3ath.com Internet Source	1 %
9	ouci.dntb.gov.ua Internet Source	1 %
10	www.researchsquare.com Internet Source	<1 %
11	Raquel Mateos, José Ricardo Pérez-Correa, Herminia Domínguez. "Bioactive Properties of Marine Phenolics", Marine Drugs, 2020 Publication	<1 %
12	e-enm.org Internet Source	<1 %

Exclude quotes On

Exclude bibliography On

Exclude assignment On
template

Exclude matches Off