

91% Unique

Total 15618 chars, 2229 words, 95 unique sentence(s).

Custom Writing Services - *Paper writing service you can trust. Your assignment is our priority! Papers ready in 3 hours! Proficient writing: top academic writers at your service 24/7! Receive a premium level paper!*

STORE YOUR DOCUMENTS IN THE CLOUD - *1GB of private storage for free on our new file hosting!*

Results	Query	Domains (original links)
Unique	They don't like a long and multiple visit treatments	-
Unique	Veneer has a transparent look and its color similar with tooth	-
Unique	There are two types of veneer, direct and indirect veneer	-
Unique	To report that veneer can correct a simple tooth malocclusion with tooth anomaly instantly	-
Unique	A 17 years old, female patient came to fix her left maxillary anterior tooth	-
Unique	Labial tooth #22 surface was prepared with round end diamond bur	-
Unique	Preparation should be incisal capping veneer type	-
Unique	A completely finished veneer should be seated on clean, dry and isolated prepared tooth	-
Unique	Veneer is a best choice to fix simple malocclusion at anterior teeth instantly	-
Unique	It has a perfect aesthetic and less visit treatment	-
Unique	Keywords : Indirect porcelain veneer, palatoversion, microdontia, instantCorrespondence : Diana Soesilo	-
Unique	Departement Conservasion, Faculty of Dentistry Hang Tuah University	-
Unique	They don't like a long and multiple visit treatment	-
Unique	A detailed family history, medical history and dental history was obtained	-

Unique	In family history, none of his family members had similar problem	-
Unique	(Figure 1)Anamnesis showed tooth #22 asymptomatic	-
Unique	Porcelain Laminates veneer for maxillary left lateral incisor was planned	-
Unique	A chamferfinish line was placed lightly subgingivally in the maxillary anterior teeth	-
7 results	The finish line should be minimum 1mm away from centric contacts	oraljournal.com ijournal.com iosrjournals.org medwinpublishers.com scribd.com es.scribd.com iosrjournals.org
Unique	The shade was selected with VITA Master shade guide	-
Unique	Patient's approval wasobtained at the time of try-in	-
Unique	The cementation appointment: Was carried out as laminate preparation and tooth preparation for cementation	-
Unique	Microdontia involving single tooth israther common condition	-
2 results	It affects most often the maxillary lateral incisor and the third molar	dentaljournals.in podj.com.pk
Unique	The term“malocclusion”wouldbemoreexpressive	-
Unique	tipped, displaced, rotated, in infra-occlusion, in supraocclusion and transposed	-
Unique	Indirect porcelain veneer has more advantages than direct composite veneer	-
Unique	To achieve esthetic andphysiologicallysoundresultsconsistently, an intraenamel preparation is usually indicated	-
Unique	Based on literature it appears that if the veneerprecision	-
Unique	This ensures minimal damage to tooth and gingiva and ensure optimal long-term prognosis	-
Unique	It has a perfect aesthetic and less visit treatment	-
11 results	Porcelain veneers can provide successful esthetic and functional long-term service for patients	pjsr.org researchgate.net researchgate.net uncch.pure.elsevier.com scribd.com es.scribd.com
Unique	Porcelain Laminate Veneers – An Esthetic Bond : Case Report	-
Unique	J Cont Med A Dent May-August 2014 Volume 2 Issue	-
Unique	Esthetic Rehabilitation by Porcelain Laminate – A Case Report	-
Unique	International Journal of Applied Dental Sciences 2015	-

6 results	Textbook of Operative Dentistry	amazon.com amazon.com en.wikipedia.org de.wikipedia.org dentistrytoday.com it.wikipedia.org
24 results	Jaypee Brothers Medical Publishers	jpmedpub.com easycounter.com amazon.com thejcdp.com e-library.net en.wikipedia.org jaypeedigital.com whsmith.co.uk amazon.com en.wikipedia.org
Unique	Ultra Thin Porcelain Laminate to Restore Esthetic of Anterior Teeth : Case Report	-
48,100 results	Shafer's Textbook of Oral Pathology	books.google.com amazon.com fcpspart1dentistry.com researchgate.net cmecde.com mebooksfree.com us.elsevierhealth.com books.google.com dencyclopedia.com arslanlibrary.com
Unique	Occlusion, Malocclusion and Method of Measurements – An Overview	-
Unique	Archives of Orofacial Sciences (2007) 2, 3-9Sherwood,	-
Unique	Essential of Operative Dentistry	-
Unique	New Delhi : Jaypee Brothers Medical Publisher	-
Unique	Sturdevant's Art and Science of Operative Dentistry	-
Unique	Ceramic Veneers : A Step – by – Step Case Report	-
Unique	Global Journal of Oral Science, 2016, 2, 20-27	-
Unique	Java 26-27 August 20163rdDentisphereP 2.39Case ReportVeneer Application As Instant Correction ForMicrodontia - Palatoversion ToothDiana SoesiloDepartment	-
Unique	Nowadays tooth aesthetic gains a lot of attention, especially for ladies who live in	-
Unique	Because of high rates of business in metropolis city, people tend to have	-
Unique	Veneer is a thin laminate restoration, which is put on labial surface of anterior	-
Unique	In this case, we used indirect porcelain veneer because it has a perfect similarity	-
Unique	Patient disturbed because she felt that the tooth has an abnormal size and asymmetrical	-
Unique	Impression is taken with rubber base impression material and sent to laboratory for veneer	-
Unique	Health Care” Shangri-la Hotel Surabaya, East Java 26-27 August 20163rdDentisphereBACKGROUNDNowadays tooth aesthetic gains a lot	-

Unique	Because of high rates of business in metropolis city, people tend to have	-
Unique	Advancements in the field of cosmetic and aesthetic dentistry have provided the dental professionals with	-
Unique	There are various ways to treat cosmetic dental problems depending upon the problem per	-
Unique	Diastema,tooth size discrepancy, discolorations, staining, fractures in teeth, endodontic treatment, and smile designing are	-
Unique	As being less invasive, for both hard and soft tissues andgranting satisfactory aesthetic outcome, the	-
Unique	In addition, the modern improvement of composite cements, adhesive systems and simplified cementation procedures	-
Unique	of the proximal surfaces of teeth requiring esthetic restoration.2Porcelain laminates has the following advantages: tooth	-
Unique	longevity than that of resincomposite veneers.3,4CASE AND CASE MANAGEMENTA 17 years old female patient came	-
Unique	Patient disturbed because she felt that the tooth has an abnormal size and asymmetrical,	-
6 results	The patient was unhappy with the appearance of her teeth and restrained herself from	oraljournal.com pjsr.org jcmad.com yumpu.com scribd.com
Unique	Clinical examination showed that tooth # 22 vital, nohypersensitivity, microntia, palatoversion, surrounded by normal	-
Unique	Proceeding Book ISBN 978-602-14590-1-0Page 338Current Concepts and Technology in Improving Dental and Oral Health	-
Unique	# 22 microdontia and palatoversionOwing to its minimally invasive nature and excellent aesthetic qualities	-
Unique	Depth orientation grooves were placed on the facial surface of the tooth with 0.3mm and	-
Unique	end tapered diamond so the prismatic top surface of mature unprepared enamel, which is known	-
Unique	Distally the tooth preparation was extended into the contact area but terminated facial to	-
Unique	An overlapped incisal edge preparation was chosen because incisal overlap provides a vertical stop that	-
Unique	The lingual finish line was placed with a round end tapered diamond, approximately one	-
Unique	The veneer extended onto thelingual surface will enhance mechanical retention and increase the surface	-
Unique	and Oral Health Care” Shangri-la Hotel Surabaya, East Java 26-27 August 20163rdDentisphereAfter gingival retraction, impression	-
Unique	Register initial tooth colorProvisional restorations were not required as the tooth reduction was minimal	-

Unique	The porcelain laminates were fabricated and were tried in for shade, fit, marginal adaptation,	-
Unique	Dual cure composite crown and bridge luting agent (Rely X, 3M, USA) was to	-
Unique	#22 Porcelain veneer insertionDISCUSSIONThe term of microdontia is used when teeth are smaller than	-
Unique	According to Shafer et al there are three types of microdontia : true generalized	-
Unique	lateral incisor, and this condition has been called the “peg lateral”.5Palatoversion is one of teeth	-
Unique	August 20163rdDentisphereteeth that were twisted or unevenly arranged, did not express the full meaning of	-
10 results	treatment “if the disfigurement or functional defect was likely to be an obstacle to the	ufrgs.br saflii.org gatesofvienna.blogspot.com gesetze-im-internet.de onelook.com corporateofficehq.com manythings.org ambrosebierce.org superheronation.com condoids.com
9 results	of individual teeth in each arch: a tooth in an arch may occupy a position	kck.usm.my scribd.com scribd.com slideshare.net readbag.com prezi.com docplayer.net es.scribd.com
Unique	like a thin layer of tooth colored material that is applied to a tooth for	-
Unique	discoloration resistant to bleaching, developmental enamel defects, closing spaces in anterior teeth, correcting mildmalalignment of	-
Unique	There are two types veneer based on method of fabrication : Direct veneers with	-
Unique	prone to fractures thanother types of veneers.3Procedure of tooth veneer preparation is cleaning the teeth,	-
Unique	The only exception is in cases in which the facial aspect of the tooth	-
Unique	In these cases, mere roughening of the involved enamel and defining of the peripheral	-
5 results	preparations that involve no reduction of the incisal edge or preparation of the lingual surfaces,	pdfs.semanticscholar.org scribd.com slideshare.net slideshare.net docplayer.net
Unique	Technology in Improving Dental and Oral Health Care” Shangri-la Hotel Surabaya, East Java 26-27 August	-
Unique	Tooth preparation that incorporates incisal edgeis preferable, because the veneer is stronger and provides	-
Unique	Moreover, this preparation design is simple and the incisal translucency is easier to becreated	-
Unique	If the gingival margins are well isolated and spaced from the retraction cord, the	-
Unique	for recording the gingival margin is afforded by cord removal just before injection of the	-
Unique	It also is recommended that the lingual aspect of the gingival embrasures be blocked	-

Unique	which often results in torn impressions, especially along criticalmarginal areas.8light-cured resin cement is recommended for	-
Unique	Attempting to achieve the color compatibility among the porcelain, tooth, and cement, the cement	-
Unique	marginal staining and their effect on gingival tissues ideally 0.3 mm of thickness for each	-
Unique	Despite following all precautions, because of the delicate nature of porcelain veneers,possible post-operative complication	-
Unique	an aesthetic concern, the patient should be informed and the veneersshould be left in place.2RESULTVeneer	-
Unique	Porcelain laminate veneer offers more extensive applications when they are used cautiously and the	-
Unique	It has become increasingly apparent that conservation of tooth structure is a major factor	-
Unique	2014 Oct-Dec;11(4):417-22Proceeding Book ISBN 978-602-14590-1-0Page 342Current Concepts and Technology in Improving Dental and	-

Top plagiarizing domains: **scribd.com** (6 matches); **amazon.com** (5 matches); **es.scribd.com** (3 matches); **slideshare.net** (3 matches); **en.wikipedia.org** (3 matches); **researchgate.net** (3 matches); **books.google.com** (2 matches); **oraljournal.com** (2 matches); **pjsr.org** (2 matches); **docplayer.net** (2 matches); **iosrjournals.org** (2 matches); **jcmad.com** (1 matches); **gatesofvienna.blogspot.com** (1 matches); **gesetze-im-internet.de** (1 matches); **saflii.org** (1 matches); **ufrgs.br** (1 matches); **yumpu.com** (1 matches); **onelook.com** (1 matches); **kck.usm.my** (1 matches); **readbag.com** (1 matches); **prezi.com** (1 matches); **pdfs.semanticscholar.org** (1 matches); **arslanlibrary.com** (1 matches); **condoids.com** (1 matches); **manythings.org** (1 matches); **ambrosebierce.org** (1 matches); **superheronation.com** (1 matches); **corporateofficehq.com** (1 matches); **fcpspart1dentistry.com** (1 matches); **de.wikipedia.org** (1 matches); **dentistrytoday.com** (1 matches); **it.wikipedia.org** (1 matches); **uncch.pure.elsevier.com** (1 matches); **podj.com.pk** (1 matches); **ijoindental.com** (1 matches); **medwinpublishers.com** (1 matches); **dentaljournals.in** (1 matches); **jpmepub.com** (1 matches); **easycounter.com** (1 matches); **cmecde.com** (1 matches); **mebooksfree.com** (1 matches); **us.elsevierhealth.com** (1 matches); **whsmith.co.uk** (1 matches); **jaypeedigital.com** (1 matches); **thejcdp.com** (1 matches); **e-library.net** (1 matches); **dencyclopedia.com** (1 matches);

Current Concepts and Technology in Improving Dental and Oral Health Care” Shangri-la Hotel Surabaya, East Java 26-27 August 20163rdDentisphereP 2.39Case ReportVeneer Application As Instant Correction ForMicrodontia - Palatoversion ToothDiana SoesiloDepartment Conservation, Fakultas of Dentistry Hang Tuah UniversityABSTRACTBackground. Nowadays tooth aesthetic gains a lot of attention, especially for ladies who live in metropolis city like Surabaya. Because of high rates of business in metropolis city, people tend to have a perfect tooth aesthetic instantly. They don't like a long and multiple visit treatments. Veneer is a thin laminate restoration, which is put on labial surface of anterior tooth. Veneer has a transparent look and its color similar with tooth. There are two types of veneer, direct and indirect veneer. In this case, we used indirect porcelain veneer because it has a perfect similarity with tooth and never going to change color. Purpose. To report that veneer can correct a simple tooth malocclusion with tooth anomaly instantly. Case. A 17 years old, female patient came to fix her left maxillary anterior tooth. Patient disturbed because she felt that the tooth has an abnormal size and asymmetrical also didn't locate in the right curve of spee. Case management. Labial tooth #22 surface was prepared with round end diamond bur. Preparation should be incisal capping veneer type. Impression is taken with rubber base impression material and sent to laboratory for veneer formation. A completely finished veneer should be seated on clean, dry and isolated prepared tooth. Conclusion. Veneer is a best choice to fix simple malocclusion at anterior teeth instantly. It has a perfect aesthetic and less visit treatment.Keywords : Indirect porcelain veneer, palatoversion, microdontia, instantCorrespondence : Diana Soesilo. Departement Conservation, Faculty of Dentistry Hang Tuah University. Arif Rahman Hakim 150. Telephone. +62 31 5912191. Email:dianasoesilo@yahoo.comProceeding Book || ISBN 978-602-14590-1-0Page 337Current Concepts and Technology in Improving Dental and Oral Health Care” Shangri-la Hotel Surabaya, East Java 26-27 August

20163rdDentisphereBACKGROUNDNowadays tooth aesthetic gains a lot of attention, especially for ladies who live in metropolis city like Surabaya. Because of high rates of business in metropolis city, people tend to have a perfect tooth aesthetic instantly. They don't like a long and multiple visit treatment.Advancements in the field of cosmetic and aesthetic dentistry have provided the dental professionals with new opportunities in conservative and aesthetic restorative procedures. There are various ways to treat cosmetic dental problems depending upon the problem per se. Diastema,tooth size discrepancy, discolorations, staining, fractures in teeth, endodontic treatment, and smile designing are some of the reasons for which patient seekaesthetic dental treatment1.As being less invasive, for both hard and soft tissues andgranting satisfactory aesthetic outcome, the rehabilitation procedure with porcelain veneers has been widely welcomed by the patients. In addition, the modern improvement of composite cements, adhesive systems and simplified cementation procedures also enable the promotion of this effective treatment approach among the dentists. Porcelain veneers is a thin bonded ceramic restoration that restores the facial surface andpart of the proximal surfaces of teeth requiring esthetic restoration.2Porcelain laminates has the following advantages: tooth structure preservation because of minimum or none prepare; thermal expansion similar to that of enamel, appearance similar to that of tooth, color stability, biocompatibility with periodontaltissues, resistance to wear, reinforcement of tooth structure, gloss retention, longer clinical longevity than that of resincomposite veneers.3.4CASE AND CASE MANAGEMENTA 17 years old female patient came to fix her left maxillary anterior tooth which was microdontia and palatoversion. Patient disturbed because she felt that the tooth has an abnormal size and asymmetrical, also didn't locate in the right curve of spee. **The patient was unhappy with the appearance of her teeth and restrained herself from** smiling due to self-consciousness. A detailed family history, medical history and dental history was obtained. In family history, none of his family members had similar problem. (Figure 1)Anamnesis showed tooth #22 asymptomatic. Clinical examination showed that tooth # 22 vital, nohypersensitivity, microntia, palatoversion, surrounded by normal gingiva.Proceeding Book || ISBN

978-602-14590-1-0Page 338Current Concepts and Technology in Improving Dental and Oral Health Care” Shangri-la Hotel Surabaya, East Java 26-27 August 20163rdDentisphereFigure 1. # 22 microdontia and palatoversionOwing to its minimally invasive nature and excellent aesthetic qualities it was decided to enhance her appearance using porcelain laminate veneers. Porcelain Laminates veneer for maxillary left lateral incisor was planned.Depth orientation grooves were placed on the facial surface of the tooth with 0.3mm and 0.5mm three wheel diamond depth cutter on the gingival half and incisal half respectively. The tooth structure remaining between the depth orientation grooves were removed with a round end tapered diamond so the prismatic top surface of mature unprepared enamel, which is known to offer only a minor retention capacity, was removed. A chamferfinish line was placed lightly subgingivally in the maxillary anterior teeth. Distally the tooth preparation was extended into the contact area but terminated facial to the contact area.An overlapped incisal edge preparation was chosen because incisal overlap provides a vertical stop that aids in the proper seating of the veneer. The lingual finish line was placed with a round end tapered diamond, approximately one fourth the way down the lingual surface connecting the two proximal finish lines. **The finish line should be minimum 1mm away from centric contacts.** The veneer extended onto thelingual surface will enhance mechanical retention and increase the surface area for bonding. (Figure 2)Figure 2. # 22 Tooth preparationProceeding Book || ISBN 978-602-14590-1-0Page 339Current Concepts and Technology in Improving Dental and Oral Health Care” Shangri-la Hotel Surabaya, East Java 26-27 August 20163rdDentisphereAfter gingival retraction, impression of maxillary arch was obtained with irreversible hydrocolloid rubber base impression material withcustomized impression tray. The shade was selected with VITA Master shade guide. (Figure 3)Figure 3. Register initial tooth colorProvisional restorations were not required as the tooth reduction was minimal and restricted to enamel. The porcelain laminates were fabricated and were tried in for shade, fit, marginal adaptation, shape, size, symmetry and contacts. Patient's approval wasobtained at the time of try-in.The cementation appointment: Was carried out as laminate preparation and tooth preparation for cementation. Dual cure composite crown and bridge luting agent (Rely X, 3M, USA) was to be used for cementation. (Figure 4)Figure 4. # 22 Porcelain veneer insertionDISCUSSIONThe term of microdontia is used when teeth are smaller than normal. According to Shafer et al there are three types of microdontia : true generalized microdontia, relativegeneralized microdontia, and microdontia involving single tooth. Microdontia involving single tooth israther common condition. **It affects most often the maxillary lateral incisor and the third molar.** One of the common forms of localized microdontia is that which affects the maxillary lateral incisor, and this condition has been called the “peg lateral”.5Palatoversion is one of teeth malocclusion conditions. The term

“irregularities of teeth” as applied toProceeding Book || ISBN 978-602-14590-1-0Page 340Current Concepts and Technology in Improving Dental and Oral Health Care” Shangri-la Hotel Surabaya, East Java 26-27 August 20163rdDentisphereTeeth that were twisted or unevenly arranged, did not express the full meaning of these deformities. The term“malocclusion”wouldbemoreexpressive. The World Health Organization (1987), had included malocclusion under the heading of Handicapping Dento Facial Anomaly, defined as an anomaly which causes disfigurement or which impedes function, and requiring **treatment “if the disfigurement or functional defect was likely to be an obstacle to the** patient's physical or emotional well-being”.Proffit (1986) elaborated that malocclusion might be associated with one or more of the following:Malalignment **of individual teeth in each arch: a tooth in an arch may occupy a position** deviating from the smooth curve of line by being; tipped, displaced, rotated, in infra-occlusion, in supra-occlusion and transposed.Malrelationship of the dental arches relative to the normal occlusion: may occur in any of the three planes of spaces: anteroposterior, vertical ortransverse.6A veneer is labial partial crown which looked like a thin layer of tooth colored material that is applied to a tooth for esthetically restoring localized or generalized defects of orintrinsic discolorations. Veneer indications are for fracture anterior teeth, large noncarious cervical lesion on anterior teeth, discoloration resistant to bleaching, developmental enamel defects, closing spaces in anterior teeth, correcting mildmalalignment of anterior teeth.3.7 There are two types of veneersaccording to design : partial veneer, full veneer. There are two types veneer based on method of fabrication : Direct veneers with composite material and indirect veneers with porcelainmaterial. Indirect porcelain veneer has more advantages than direct composite veneer. It less technique sensitive than direct veneer, multiple teeth can be done in less time, chair time required for indirect veneer is less, produce better contour, contact and shades, has longer life than direct veneer, better retention, less prone to stains, good aesthetics, less prone to fractures thanother types of veneers.3Procedure of tooth veneer preparation is cleaning the teeth, selection the shade, tooth isolation and retraction of gingiva using retraction cord. To achieve esthetic andphysiologicallysoundresultsconsistently, an intraenamel preparation is usually indicated. The only exception is in cases in which the facial aspect of the tooth is significantly undercontoured because of severe abrasion or erosion. In these cases, mere roughening of the involved enamel and defining of the peripheral margins are indicated. Intraenamel preparation (or the roughening of the surface in undercontoured areas) before placing a veneer is strongly recommended for the following reasons: to provide space for opaque, bonding, or veneering materials formaximal esthetics without overcontouring, to remove the outer, fluoride-rich layer of enamel that may be more resistant to acid etching, to create a rough surface for improved bonding, to establish a definite finish line.3.8Different preparation designs have been advocated from feather and window **preparations that involve no reduction of the incisal edge or preparation of the lingual surfaces,** to other preparations that involve reduction of the incisal edges.

Incisal preparation is carried over the incisalProceeding Book || ISBN 978-602-14590-1-0Page 341Current Concepts and Technology in Improving Dental and Oral Health Care” Shangri-la Hotel Surabaya, East Java 26-27 August 20163rdDentisphereedge from buccal to palatal, with up to 1.5-2 mm of incisal reduction. Tooth preparation that incorporates incisal edgeis preferable, because the veneer is stronger and provides a positive seat during cementation. Moreover, this preparation design is simple and the incisal translucency is easier to becreated by the ceramist.9An elastomeric impression is made of the preparations. If the gingival margins are well isolated and spaced from the retraction cord, the cord may be left in place during impression making. If the margins are subgingival or close to the gingival tissue, however, better access for recording the gingival margin is afforded by cord removal just before injection of the impression material. It also is recommended that the lingual aspect of the gingival embrasures be blocked out with soft wax if thegingival embrasures are open faciolingually. This step prevents penetration and interlocking of the impression material through the gingival embrasure, which often results in torn impressions, especially along criticalmarginal areas.8light-cured resin cement is recommended for bonding the veneer to the tooth. Attempting to achieve the color compatibility among the porcelain, tooth, and cement, the cement shade was chosen. This was achieved through shade guides and hydrosoluble pastes whose shades are equal to those of the cements and enable the evaluation of the color ontothe tooth.4On the other hand long-term study of porcelain veneers is required in order to study their marginal integrity, marginal staining and their effect on gingival tissues ideally 0.3 mm of thickness for each shade change. Based on literature it appears that if the veneerprecision. This ensures minimal damage to tooth and gingiva and ensure optimal long-term prognosis. Despite following all precautions, because of the delicate nature of porcelain veneers,possible post-operative complication is cracking. If the veneer has been well bonded to the underlying enamel and is not an aesthetic concern, the patient should be informed and the veneersshould be left in place.2RESULTVeneer is a best choice to fix simple malocclusion at anterior teeth instantly. It has a perfect aesthetic and less visit treatment. **Porcelain veneers can provide successful esthetic and functional long-term service for patients.** Porcelain laminate veneer offers more extensive applications when they are used cautiously and the results achieved have been gratifying for the cosmetic dentist and the patient alike. It has become increasingly apparent that conservation of tooth structure is a major factor in determining the long-term prognosis of any restorative procedure.REFERENCETabassum, R. 2014. Porcelain Laminate Veneers – An Esthetic Bond : Case Report. J Cont Med A Dent May-August 2014 Volume 2 Issue 2.Soni, R., Vivek, R. 2015. Esthetic Rehabilitation by Porcelain Laminate – A Case Report. International Journal of Applied Dental Sciences 2015; 1(4): 98-100Garg, N. and Garg.A. 2013. **Textbook of Operative Dentistry.** New Delhi : 2nd Ed. **Jaypee Brothers Medical Publishers.** p. 302-316Bizio, O., et.al. 2014. Ultra Thin Porcelain Laminate to Restore Esthetic of Anterior Teeth : Case Report. RSBO. 2014 Oct-Dec;11(4):417-22Proceeding Book || ISBN 978-602-14590-1-0Page 342Current Concepts and Technology in Improving Dental and Oral Health Care” Shangri-la Hotel Surabaya, East Java 26-27 August 20163rdDentisphereShafer, W.G, Hine, M.K., Levy, B.M. 2009. **Shafer's Textbook of Oral Pathology.** 6th Ed. Philadelphia: W.B. Saunders. p. 37-38Hassan, R. and Rahimah, A.K. 2007. Occlusion, Malocclusion and Method of Measurements – An Overview. Archives of Orofacial Sciences (2007) 2, 3-9Sherwood, I.A. 2010. Essential of Operative Dentistry. 1t Ed. New Delhi : Jaypee Brothers Medical Publisher. p. 451-469Heymann, H.O; Swift, E.J; Ritter, A.V. 2013. Sturdevant's Art and Science ofOperative Dentistry. 6th Ed. St.Louis : Elsevier Mosby. p. 624-688.Mancini, M. and Mancini, M. 2016. Ceramic Veneers : A Step – by – Step Case Report. Global Journal of Oral Science, 2016, 2, 20-27.