

PAPER • OPEN ACCESS

## The 12th Congress of Indonesian Society for Biochemistry and Molecular Biology in Conjunction With The 2nd International Conference “Collaboration Seminar of Chemistry and Industry (CoSCI)” and AnMicro Workshop

To cite this article: 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **217** 011001

View the [article online](#) for updates and enhancements.

## Preface

Second International Conference “Collaboration Seminar of Chemistry and Industry (CoSCI 2018) in conjunction with 23<sup>rd</sup> Indonesian Society for Biochemistry and Molecular Biology (ISBMB) seminar was taken place at Universitas Airlangga, Surabaya-Indonesia on October 11-12, 2018. The event presented a theme” Recent Development of Omics Technology For Human Prosperity”.

"Omics is a general term for a broad discipline of science and engineering for analyzing the interactions of biological information objects in organism. The main focus is on: 1) mapping information objects such as genes, proteins, and ligands; 2) finding interaction relationships among the objects; 3) engineering the networks and objects to understand and manipulate the regulatory mechanisms; and 4) integrating various omes and omics subfields.

The event was held to facilitate for the scientists, scholars, engineers and students from universities, research institutes and industries to present ongoing research activities, especially in Biochemistry, Biology, Chemistry, Medicine and other in related fields. It was also to encourage future collaboration among all participants.

The conference of CoSCI and ISBMB created proceedings from the papers that were submitted by participants, after they were reviewed by committee members and international reviewers. This volume intends to provide readers with the recent advances in the Omics Technology such as Chemistry, Biochemistry and Molecular Biology, and Medicine field.

We would like to thank to all authors who contributed to the proceedings and also to the organizing committee, reviewers, speakers, sponsor, and all the conference participants for their supporting in the conference of CoSCI 2018 and 23<sup>rd</sup> Seminar of ISBMB.

Dr. Purkan, M.Si

Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga,  
Surabaya, Indonesia, Nov. 10, 2018



PAPER • OPEN ACCESS

## Committee

To cite this article: 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **217** 011002

View the [article online](#) for updates and enhancements.

## COMMITTEE

### The Chairman

1. Dr. Purkan, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia.
2. Dr. Sri Sumarsih, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia

### Steering Committee

1. Dr. Abdulloh, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
2. Dr. Alfinda Novi Kristanti, DEA (Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
3. Dr. Ni'matuzarroh, M.Si, Biology Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
4. Satya Candra Wibawa Sakti, M.Sc, Ph.D, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
5. Harsasi Setyawati, S.Si, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
6. Ahmadi Jaya Permana, S.Si, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
7. Kautsar UI Haq, S.Si, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
8. Dr. Rico Ramadhan, M.Sc, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
9. Dr. Indeswati Diyatri, drg., M.S, Faculty of Dentistry, Universitas Airlangga
10. Dr. Dian Mulawarmanti, drg., M.S, Faculty of Dentistry, Hang Tuah University, Surabaya Indonesia

### Scientific Committee

#### Editors:

1. Ali Rohman, M.Si, Ph.D, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
2. M. Zakki Fahmi, M.Si, Ph.D, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
3. Dr. Verawat Champreda, M.Sc, National Center for Genetic Engineering and Biotechnology (BIOTEC), Thailand
4. Dr. Sehanat Prasongsuk, Department of Botany, Chulangkorn Universty, Thailand



**Reviewers :**

1. Prof. Ni Nyoman Tri Puspaningsih, M.Si, Biochemistry Research Division, Faculty of Science and Technology, Universitas Airlangga, Indonesia
2. Prof. dr. Soetjipto, MS., Ph.D, Biochemistry Department, Faculty of Medicine, Universitas Airlangga
3. Prof. Seung Wook Kim, Dept of Chemical and Biological Eng, Korea University
4. Prof. Rendy Kartika, Chemistry Department, Louisiana State University, Baton Rouge, LA
5. Prof. Hunsa Punnapayak, Department of Botany, Chulalongkorn University, Thailand
6. Prof. Antonius Suwanto, Division of Microbiology, Institut Pertanian Bogor, Indonesia
7. Prof. Dessy Natalia, Biochemistry Research Division, Faculty of Mathematics and Natural Sciences, Institut Teknologi Bandung, Indonesia
8. Prof. Dr, Suraini Abdul Aziz, Faculty of Biotechnology and Biomolecular Sciences, University Putra Malaysia
9. Prof. Afaf Baktir, Biochemistry Research Division, Faculty of Science and Technology, Universitas Airlangga, Indonesia
10. Dr. Mulyadi Tanjung, M.Si, Natural Product Research Division, Faculty of Science and Technology, Universitas Airlangga, Indonesia
11. Tjitjik Srie Tjahjandarie, Ph.D, Natural Product Research Division, Faculty of Science and Technology, Universitas Airlangga, Indonesia
12. Prof Fedik Rantam, Stem Cell Research Division, Faculty of Veterinary, Universitas Airlangga, Indonesia
13. Prof. Yosephine Sri Wulan Manuhara, Biology Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
14. Dr. Hery Suwito, M.Si (Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
15. Dr. Nanik Siti Aminah, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
16. Dr. rer. nat. Ganden Supriyanto, M.,Sc, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia
17. Dr. Pratiwi Pudjiastuti, M.Si, Chemistry Department, Faculty of Science and Technology, Universitas Airlangga, Indonesia

PAPER • OPEN ACCESS

## Peer review statement

To cite this article: 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **217** 011004

View the [article online](#) for updates and enhancements.

## Peer review statement


All papers published in this volume of *IOP Conference Series: Earth and Environmental Science* have been peer reviewed through processes administered by the proceedings Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.





# IOP Conference Series: Earth and Environmental Science

18

H Index

Country	United Kingdom -  SIR Ranking of United Kingdom
Subject Area and Category	<a href="#">Earth and Planetary Sciences</a> <a href="#">Earth and Planetary Sciences (miscellaneous)</a> <a href="#">Environmental Science</a> <a href="#">Environmental Science (miscellaneous)</a>
Publisher	<a href="#">IOP Publishing Ltd.</a>
Publication type	Conferences and Proceedings
ISSN	17551315, 17551307
Coverage	2010-2020
Scope	The open access IOP Conference Series: Earth and Environmental Science (EES) provides a fast, versatile and cost-effective proceedings publication service.
	<a href="#">Homepage</a> <a href="#">How to publish in this journal</a> <a href="#">Contact</a>



IOP Conference Series:  
Earth and Environmental...

Not yet assigned  
quartile

SJR 2019  
0.18

powered by scimagojr.com

← Show this widget in  
your own website

Just copy the code below  
and paste within your html  
code:

`<a href="https://www.scimaq`



## IOP Conference Series: Earth and Environmental Science

### Table of contents

#### Volume 217

2019

◀ Previous issue Next issue ▶

The 12th Congress of Indonesian Soc. for Biochemistry and Molecular Biology in Conjunction With The 2nd Int. Conf. "Collaboration Seminar of Chemistry and Industry (CoSCI)" and AnMicro Workshop 11-12 October 2018, Universitas Airlangga, Indonesia

Accepted papers received: 21 November 2018

Published online: 09 January 2019

[Open all abstracts](#)

**OPEN ACCESS** 011001

The 12th Congress of Indonesian Society for Biochemistry and Molecular Biology in Conjunction With The 2nd International Conference "Collaboration Seminar of Chemistry and Industry (CoSCI)" and AnMicro Workshop

◀ Open abstract View article PDF

**OPEN ACCESS** 011002

Committee

◀ Open abstract View article PDF

**OPEN ACCESS** 011003

Conference Photographs

◀ Open abstract View article PDF

**OPEN ACCESS** 011004

Peer review statement

◀ Open abstract View article PDF

### Papers

#### Chemistry

**OPEN ACCESS** 012001

Facile Sol-Gel Synthesis of Calcium Phosphates: Influence of Ca/P Ratio and Calcination Temperature

A J Permana, A T Utami, U S Handayani and H Setyawati

◀ Open abstract View article PDF

**OPEN ACCESS** 012002

Determination of Brønsted Acid Sites In Porous Aluminosilicate Solid Catalysts Using Volumetric And Potentiometric Titration Method

A Purwaningsih, A N Kristanti, D Z Mardho, D W Saraswati, N M Putri, N H Saputri and Hartati

◀ Open abstract View article PDF

**OPEN ACCESS** 012003

Carbon Paste Electrode Modified Imprinted Zeolite as a Selective Sensor for Creatine Analysis by Potentiometry

A. Athiroh, T Fadillah, D F Damayanti, A A Widati, A Abdulloh and M Khasanah

◀ Open abstract View article PDF

**OPEN ACCESS** 012004

Voltammetric Study of Ascorbic Acid Using Polymelamine/Gold Nanoparticle Modified Carbon Paste Electrode

A N Fajida, E Fitriany, A Bakhtir, F Kurniawan and M Harsini

◀ Open abstract View article PDF

**OPEN ACCESS** 012005

Synthesis of Silver Nanoparticles and the Development in Analysis Method

H I Badri'ah, F Seedeh, G Supriyanto and A H Zaidan

◀ Open abstract View article PDF

### JOURNAL LINKS

[Journal home](#)  
[Information for organizers](#)  
[Information for authors](#)  
[Search for published proceedings](#)  
[Contact us](#)  
[Reprint services from Curran Associates](#)



Activate Windows

Activate Windows

## OPEN ACCESS

012006

Two Flavonoids From Stem Bark of *Casimiroa edulis* and Their Antidiabetic and Antioxidant Activities

K N W Tun, N S Aminah, A N Kristanti, R Ramadhan and Y Takaya

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012007

## Graphene Oxide from Bagasse/Magnetite Composite: Preparation and Characterization

M Jannatin, G Supriyanto, Abdulloh, W A W Ibrahim and N K Rukman

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012008

GO-Fe<sub>3</sub>O<sub>4</sub> Nanocomposite from coconut shell: Synthesis and characterization

N K Rukman, M Jannatin, G Supriyanto, M Z Fahmi and W A W Ibrahim

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012009

## First Order Kinetics of Salicylamide Release from κ-Carrageenan Hard Shell Capsules in Comparison with Gelatin

P Pudjastuti, E Hendradi, S Wafiroh, H Darmokoeseomo, M A R D Fauzi, L Nahar and S D Sarker

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012010

Chromanone Acid Derivatives from the Stem Bark of *Calophyllum incassatum*

U Hasanah, T S Tjahjandarie and M Tanjung

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012011

Preparation Hydrophobic Fabric Coated by TiO<sub>2</sub> and Hexadecyltrimethoxysilane

U S Handayani, A A Widati and I N Yusbainika

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012012

Kecombrang (*Etlingera elatior*) Leaves Ethanol Extract Effect to Lens and Erythrocyte Aldose Reductase Activity in Wistar strain white rats (*Rattus norvegicus*) Streptozotocin induced

S Handayani, H Notopuro and G I Prabowo

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012013

## Adsorption of Isopropyl Alcohol (IPA) in Water Using Activated Bentonite

A Abdulloh, G Supriyanto and O W Ningsih

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012014

## Production of Nanopropolis Using High Pressure Ball Mill Homogenizer

D Hamdi, A Wijanarko, H Hermansyah, S C Asih and M Sahlan

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012015

Synthesis of ZnO-TiO<sub>2</sub>/Chitosan Nanorods By Using Precipitation Methods and Studying Their Structures and Optics Properties at Different Precursor Molar Compositions

Y Rilda, D Damara, Syukri, Y E Putri, Refinel and A Agustien

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012016

Phytochemical Screening and Antioxidant Activity of Ethanol Extract of Leilem (*Clerodendrum minahassae* Teijsm. & Binn) as an Antihyperlipidemic and Antiatherosclerotic Agent

C F Kalirupan, F R Mantiri and R R H Rumende

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012017

## Concentration of Some Metals in Water and Soil Samples at Some Locations near the Hotmud Flow at Porong Disaster Area, Sidoarjo, East Java, Indonesia.

A Wiryawan, R Suntari, Z Kusuma and Syekhtani

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012018

The Effect of Roselle (*Hibiscus sabdariffa* Linn) Flower Extract To The SGPT Activity In Male Wistar Rats (*Rattus Norvegicus*) Induced By High Dose Paracetamol

D Halim, E J Sihning and Tehupuring

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012019

## Antioxidant Exploration in Cardamom Rhizome Potential as a Functional Food Ingredient

H Winarsi, A Yuniaty and Warsinah

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012020

Effect of Gambir Catechin Isolate (*Uncaria Gambir* Roxb.) Against Rat Triacylglycerol Level (*Rattus novergicus*)

Y Alioes, R R Sukma and S L Sekar

[+ Open abstract](#) [View article](#) [PDF](#)

## Biochemistry and Molecular Biology

## OPEN ACCESS

012021

## Exploration of Cellulolytic Microorganism as A Biocatalyst Candidate for Liquid Fertilizer Production

N Halimah, A Bakir and P Purkan

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012022

Antibody Titers in The Sheep which were Immunated Antigen of *Whole* Protein from Third Instar Larvae *Musca domestica*

B Ariantini, H Ratnani, E M Luqman and P Hastutiek

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012023

Lemon (*Citrus limon*) Juice Has Antibacterial Potential against Diarrhea-Causing Pathogen

ER Ekawati and W Darmanto

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012024

Genetic Relationship of *Hibiscus* spp. Based on DNA bands Using RAPD Technique

Hamidah and A Z Muhtadi

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012025

Effect of *Sticophus hermanni* extract on fasting blood glucose and skeletal muscle glut4 on type 2 diabetes mellitus rats model

I Safitri, B Purwanto, L Rochyani, G I Prabowo and D Sukmaya

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012026

Callus Induction and Bioactive Compounds from *Piper betle* L. var *nigra*

Junairiah, A Mahmuda, Y S W Manuhara, Ni'matuzahroh and L Sulistyorini

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012027

Antimicrobial Activity of Ethanol Extract of *Abrus precatorius* L. Roots against Planktonic Cells and Biofilm of Urine and Blood Methicillin Sensitive *Staphylococcus aureus* (MSSA) Isolate

B Mutmalnah, Ni'matuzahroh and A Bakir

## OPEN ACCESS

012028

Utilization of Rice Straw Hydrolysis Product of *Penicillium* sp. H9 as A Substrate of Biosurfactant Production by LI161 Hydrocarbonoclastic Bacteria

Ni'matuzahroh, S K Sari, N Trikurniadewi, A D Pusfita, I P Ningrum, S N M M Ibrahim, T Nurhariyati, Fatimah and T Surtiningsih

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012029

Carbon and Nitrogen Sources for Lipase Production of *Micrococcus* sp. Isolated from Palm Oil Mill Effluent-Contaminated Soil

S. Sumarsih, S. Hadi, D.G.T. Andini and F.K. Natsihana

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012030

Cytotoxicity of Combination Chitosan with Different Molecular Weight and Ethanol Extracted *Aloe vera* using MTT Assay

Sularsih, Soetjipto and Retno Pudji Rahayu

[+ Open abstract](#) [View article](#) [PDF](#)

## OPEN ACCESS

012031

## Hepatoprotective Effect of Gamma-mangostin for Amelioration of Impaired Liver Structure and Function in Streptozotocin-induced Diabetic Mice

S A Husen, D Winarni, Salamun, A N M Anson, R J K Susilo and S Hayaza

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012032

Utility of *Saccharomyces cerevisiae* As Probiotics to Induce Protease Production For Worms Feed Improvement

R Arissirajudin, S Hadi, Abdillan Safa and P Purkan

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012033

Induction of Angiogenesis Process in Mandible Using *Anadara granosa* Shell Graft (Experimental Laboratory Study on *Rattus norvegicus*)

Widyastuti M Rubianto and Soetjipto

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012034

Dehalogenase enzyme activity of *Bacillus* sp. D1 isolated from pharmaceutical waste

K Primasari, D W Sawitri, R Fikri, N Trikumadewi, Ni'matuzahroh and G Supriyanto

[+ Open abstract](#)

[View article](#)

[PDF](#)

OPEN ACCESS

012035

The impact of conditioned medium of umbilical cord-derived mesenchymal stem cells toward apoptosis and proliferation of glioblastoma multiforme cells

Novi Silvia Hardiany, Yohana and Septelia Inawati Wanandi

[+ Open abstract](#)

[View article](#)

[PDF](#)





UNIVERSITAS AIRLANGGA



AIRLANGGA GLOBAL ENGAGEMENT



AnMicro

ASEAN Network  
on Microbial Utilization



## CERTIFICATE FOR ATTENDANCE

*This certificate is awarded to*

**Sularsih**

*Who has participated in the 2<sup>nd</sup> International Conference "Collaboration Seminar of Chemistry and Industry (CoSCI) as  
Poster Presenter*

Held at Universitas Airlangga, Campus C Surabaya, Indonesia, October 11<sup>th</sup> – 12<sup>th</sup> October 2018

Surabaya, October 12<sup>th</sup> 2018

**Dr. Purkan, M.Si.**

Chairman of Organizing Committee

**Prof. Dr. Nyoman Tri Puspaningsih, M.Si.**

Chairman of 12<sup>th</sup> Congress and 23<sup>rd</sup> Seminar of ISBMB  
AnMicro and Marine Workshop



UNIVERSITAS AIRLANGGA



**AnMicro**  
ASEAN Network  
on Microbial Utilization



## CERTIFICATE FOR ATTENDANCE

*This certificate is awarded to*

**Sularsih**

*Who has participated in the 23<sup>rd</sup> Seminar of Indonesian Society for Biochemistry and Molecular Biology as*  
**Poster Presenter**

Held at Universitas Airlangga, Campus C Surabaya, Indonesia, October 11<sup>th</sup> – 12<sup>th</sup> October 2018

Surabaya, October 12<sup>th</sup> 2018

Dr. Rahmawati Ridwan, Apt., MS.  
President of ISBMB

Prof. Dr. Nyoman Tri Puspaningsih, M.Si.  
Chairman of 12<sup>th</sup> Congress and 23<sup>rd</sup> Seminar of ISBMB  
AnMicro and Marine Workshop

PAPER • OPEN ACCESS

## Cytotoxicity of Combination Chitosan with Different Molecular Weight and Ethanol Extracted *Aloe vera* using MTT Assay

To cite this article: Sularsih *et al* 2018 *IOP Conf. Ser.: Earth Environ. Sci.* **217** 012030

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.



# Cytotoxicity of Combination Chitosan with Different Molecular Weight and Ethanol Extracted *Aloe vera* using MTT Assay

Sularsih<sup>1,2,\*</sup>, Soetjipto<sup>3</sup>, Retno Pudji Rahayu<sup>4</sup>

<sup>1</sup>Doctoral Study Program, Faculty of Medicine, Universitas Airlangga Surabaya, Indonesia,

<sup>2</sup>Departement of Dental Material, Faculty of Dentistry, Universitas Hang Tuah, Surabaya, Indonesia,

<sup>3</sup>Department of Biochemistry, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia,

<sup>4</sup>Department of Oral Pathology Anatomy, Faculty of Dentistry, Universitas Airlangga, Surabaya, Indonesia,

\*corresponding author :larsihdentist@gmail.com

**Abstract.** The use of combination chitosan and ethanol extracted *Aloe vera* is considered to have activity to promote bone healing. The requirement dental material used in oral cavity include good biocompatibility, non toxic, non carcinogenic and nor allergenic response. The aim of this study was to examine the cytotoxicity combination chitosan with different molecular weight and ethanol extracted *Aloe vera* using MTT assay. 1 % chitosan gel (w/v) was made from chitosan powder with high (ChH) and low (ChL) molecular weight. 1 % chitosan gel was combined with ethanol extracted *Aloe vera* with concentration 50 % (Av<sub>1</sub>) and 75% (Av<sub>2</sub>). It divided five groups: control group, ChH-Av<sub>1</sub>, ChH-Av<sub>2</sub>, ChL-Av<sub>1</sub> and ChL-Av<sub>2</sub>. Each of sample was immersed in eppendorf microtubes consist of media culture. After 24 hours, the immersion of media culture were used to investigate the cytotoxic effect to BHK-21 cell lines by MTT assay method. The density of optic formazan was indicated the number of living cells. All data were statistically analyzed by one-way Anova. The result showed that percentage of the living cells of all groups was uper 50 % using parameter CD<sub>50</sub>. There was no significant difference among treatment groups. It can be concluded that combination chitosan with different molecular weight and ethanol extracted *Aloe vera* were non toxic for BHK-21 cell culture.

**Keyword :** Cytotoxicity, Chitosan, Ethanol, *Aloe vera*, MTT Assay

## 1. Introduction

Lidah buaya (Indonesian) or *Aloe vera* is natural herbal plant that easy grow in Batu, Malang, East Java, Indonesia. Batu's climate is classified as tropical. The average temperature in Batu is 21.4 °C and average annual rainfall is 3416 mm.[1] *Aloe vera* is called “the plants immortality” are proven use as herbal remedy in dentistry as anti inflammatory, antiviral, antibacteria and antioxidative.[2] It contain anthraquinones components namely aloin, aloe emodin, and barbaloin, carbohydrate, acetyl mannan glukomannan, alkaline phosphatase, bradikinin enzymes, healing hormones auxin, gibberellin, saponins and protein lectin that can enhance tissue and bone healing proces.[2; 3]

In the present dentistry study, *Aloe vera* stimulating PDLC proliferation, expression of Runx2, GDF-5, BMP-2, VEGF, type I collagen, ALPase activity, and mineral deposition. Runx2 is a transcription factor considered to be a regulator of osteoblast/cementoblast differentiation and function.[4] BMP-2, GDF-5 and VEGF are important growth factors for periodontal tissue and bone healing. Ethanol extracts *Aloe vera* have strongest antibacteria and antioxidative.[5; 6]

Chitosan is a linear polysaccharide derived from partial deacetylation of chitin. Chitosan have great utility because of their biological activities such as biocompatibility, biodegradability, non-toxicity and adsorptive abilities, as well as chemical applications, mainly in the medical and pharmaceutical fields. The characteristic properties of chitosan are related to its weight averaged molecular weight





(MW).[7] In vitro biological performance combination chitosan and *Aloe vera* offer good environment for hDF attachment, proliferation and viability. It has good cellular response, suitable mechanical and physical properties.[8]. In our previous study, the application of gel combination chitosan and ethanol extracted *Aloe vera* both concentration 50 % and 75 % increased fibroblast proliferation and collagen type 1 on wound healing process of traumatic ulcer.

The requirement dental material used in oral cavity include good biocompatibility, non toxic, non carcinogenic and non allergenic response. Therefore toxicity testing is needed.[9; 10]. The aim of this study is to examine the cytotoxicity combination chitosan with different molecular weight and ethanol extracted *Aloe vera* using MTT assay.

## 2. Experimental Method

The leaves of the plant *Aloe vera* L obtained from Batu, Malang, Jawa timur, Indonesia. Chitosan powder purchased from Sigma chemical, St. Louis, USA. The degree of deacetylation was more than 75 %. Chitosan powder with high molecular weight, 310.000 – 375.000 Da and chitosan powder with low molecular weight 50.000 – 190.000 Da.

The dried *Aloe vera* leaves were finely chopped and then soaked with 70% ethanol solvent in the macerator for  $2 \times 24$  h. The macerate was thickened with a rotary evaporator and evaporated over a water bath to remove the residual solvent to obtain a thick viscous extract with a constant weight. This research using ethanol extracts *Aloe vera* with concentration was 50 % and 75 %. Chitosan gel was made from chitosan powder with high (ChH) and low (ChL) molecular weight. 1% chitosan gel (w/v) was made with diluted one gram of chitosan powder in 100 ml 2 % acetic acid. It added with NaOH solution to get neutral pH. The mixture was stirred until the gel was completely formed. 1% chitosan gel was combined with ethanol extracted *Aloe vera* with concentration 50 % (Av<sub>1</sub>) and 75% (Av<sub>2</sub>). Combination chitosan and extracted ethanol *Aloe vera* was made with ratio 50:50.

The sample divided five groups: control group, the combination of ChH-Av<sub>1</sub>, ChH-Av<sub>2</sub>, ChL-Av<sub>1</sub> and ChL-Av<sub>2</sub>. The sample was then ready to be tested with cell culture. The control group was that each well plate had to be contained only with cells and culture media. BHK-21 cell culture in cell-line form was cultivated inside a bottle. After confluent, the cell culture was harvested by using trypsin versene solution. The harvesting result was taken out little bit and then cultivated again into Rosewell Park Memorial Institute (RPMI-1640) media containing with 10% bovine serum albumin incubated for 24 hours with temperature at 37° C. Those cells, afterwards, were put into small bottles with  $2 \times 10^5$  cell/ml density as the testing samples. This toxicity testing used 96 well plates of cell culture with flat base (Figure 1). The testing was done based on standard protocol required for MTT assay. Each plate containing cells with  $2 \times 10^5$  densities into 100 µl culture media. Before being tested, those samples had to be sterilized by ultra violet rays for 15 minutes. The samples were then put into well plates about 50 µl. Those well plates were then incubated for 20 hours with temperature at 37° C. After that, each plate was contained with 5 mg/ml MTT which had been diluted in 25 ml PBS, and those well plates were then incubated for 4 hours with temperature at 37° C. The next step, samples were taken from each well plate which was added with 50 µl DMSO and piped up and down in order to dilute crystals formed. Those well plates were then incubated for 5 minutes with temperature at 37° C. Those well plates was monitored by ELISA reader with 630 λ then shown in optical density (absorbents). The amount of absorbents in each well plate showed the number of viability cell in media culture. In order to analyze the percentage of living cell, the following formulation could be used. The percentage of the living cells that indicator it not toxic up to 50 % using parameter CD<sub>50</sub>. [9,10]

$$\% \text{ living cell} = \frac{\text{Treatment} + \text{Media}}{\text{cell} + \text{Media}} \times 100\%$$



**Figure 1.** Cytotoxicity test on 96 well plates that was monitored by ELISA reader


The data result was then tabulated based on each groups, and tested by One-Way ANOVA with 5% level of significance. If there was significant difference, LSD test would be done.

### 3. Result and Discussion

The result of cytotoxicity testing of combination chitosan with different molecular weight and ethanol extracted *Aloe vera* on cell culture in BHK-21 evaluated depends on the number of living or viability cell shown as optical density can be shown in table 1. The graphic of optic density value and viability cell can be shown in figure 2. Figure 3 shown the viability cell of each group.

**Table 1.** Average formazan optic density value, percentage of living cell and standar deviation

Group	Optic Density value $\bar{X} \pm SD$	Viability cell (%) $\bar{X} \pm SD$	n
Control	0,503 $\pm$ 0,026	100 $\pm$ 0,00	8
ChH-Av1	0,513 $\pm$ 0,034	93,087 $\pm$ 1,903	8
ChH-Av2	0,426 $\pm$ 0,038	91,643 $\pm$ 5,006	8
ChL-Av1	0,376 $\pm$ 0,098	88,568 $\pm$ 6,533	8
ChL-Av1	0,386 $\pm$ 0,017	88,05 $\pm$ 4,461	8



**Figure 2.** The graphic of optic density value and viability cell of control group, the combination of ChH-Av<sub>1</sub>, ChH-Av<sub>2</sub>, ChL-Av<sub>1</sub> and ChL-Av<sub>2</sub>



**Figure 3.** The viability cell of control group (a), the combination of ChH-Av<sub>1</sub> (b), ChH-Av<sub>2</sub> (c), ChL-Av<sub>1</sub> (d) and ChL-Av<sub>2</sub> (e)

Based on table 1, the formazan optic density values of Ch-Av from chitosan with high molecular weight more higher than Ch-Av from chitosan with low molecular weight. The percentage of the

living cells of all groups was uper 50 % using parameter  $CD_{50}$ . Kolmogorof Smirnof Test shows that all groups have probability values more than 0.05 ( $p > 0.05$ ). It means that all groups have normal distribution. In homogeneity test, however, the value obtained is about 0.130 ( $p > 0.05$ ) which means that all groups are homogeny. Thus, one way Anova test was needed to be done. The result of One-Way ANOVA test shows that there is no significant difference of optic density and viability cell among treatment group with probability value ( $p > 0.05$ ). The Combination chitosan with different molecular weight and ethanol extracted *Aloe vera* were non toxic for BHK-21 cell culture.

Cytotoxicity assays are based on the use of various parameters quantifying cell death or measuring the effect of the product on cell metabolism. This research using BHK-21 cell culture in cell-line by MTT assay method to measure the cytotoxicity combination chitosan with different molecular weight and ethanol extracted *Aloe vera*. BHK-21 cell from fibroblast of baby hamster's kidney is often used by the researchers for cytotoxicity testing of dentistry materials since it is the most important cell in the components of pulp, ligament periodontal and gingiva.[11] MTT assay is the most common methods used to determine cytotoxicity. The test was proved to be more accurate and time saving than other conventional haemocytometer counting methods. The well with the highest absorbance indicates the highest viability in the case of MTT assay.[12] The MTT assay is the most used in vitro method for determining cytotoxicity by measuring cell metabolism, using a reaction catalyzed by the succinate dehydrogenase enzyme from mitochondria of viable cells. Trypan blue is an assay that allows differentiation viable cells by microscopic evaluation, since the cells that present loss of membrane integrity allow the incorporation of this agent into the nucleus. The reparative materials evaluated by MTT and Trypan blue cytotoxicity assays demonstrated viability in all evaluated cell lines.[13]

The percentage of the living cells of all groups in this research was uper 50 % using parameter  $CD_{50}$ . There was no significant difference among treatment groups. It means that combination chitosan with different molecular weight and ethanol extracts *Aloe vera* were non toxic for BHK-21 cell culture. There is no toxicity on BHK-21 cell culture since chitosan does not have function and structure which can make chitosan and *Aloe vera* becomes toxic. The process of chitosan production, moreover, has passed deproteination phase which is a phase of separating or breaking bond between protein and chitin so that chitosan will not become toxic.[9] Combination chitosan with high molecular weight and ethanol extracted *Aloe vera* have high average formazan optic density value than combination chitosan with low molecular weight and ethanol extracted *Aloe vera*. The principal of MTT assay is to break tetrazolium MTT ring, (3-(4,5-dimethylthiazol-2-yl)-2,5diphenyl tetrazolium bromide), which has yellow color because of dehydrogenesis in active mitochondria, and to produce insoluble blue-purple formazan product. If the dehydrogenises are not active because of cytotoxic effects, the formazan is not produced. The formazan production can be measured by dissolving it and measuring optic density of the solution produced. The lower the percentage of optic density is, the fewer the number of active metabolic cells that can reduce MTT.[9; 14] The biological properties of chitosan depend on their physicochemical parameters, especially their solubility. The characteristic of chitosan is related with its molecular weight.[15] Chitosan with high molecular weight powder become chitosan gel with good physical characteristic and higher viscosity. Powder with high molecular weight has greater particle size, due to the larger size of the solute molecules solvent molecules surrounding more increasingly difficult to dissolve. It become tenuous bonds by the force of molecular attraction.[16] The molecular weight is also directly related to length of molecular chain, more longer the molecular chain, the viscosity more higher and more higher average formazan optic density value.[14; 16]

The result of this study, combination chitosan and ethanol extracted *Aloe vera* was not toxic. Chitosan is used in various biomedical applications due to its significant antibacterial activity, biodegradability, non toxicity, and biocompatibility. The synergistic association combination of chitosan and ethanol extracts *Aloe vera* have biological performance that offer good enviroment for viability of fibroblast cell. It regarding requirement for wound healing such as control inflammation, immunomodulatory activity, epithelialization and fibroblast proliferation.[17; 18]

#### 4. Conclusion

Based on this research, the percentage of the living cells of group combination chitosan and ethanol extracted *Aloe vera* was up to 50 %. It shown that the combination chitosan with different molecular weight and ethanol extracted *Aloe vera* were non toxic with BHK-21 cell lines by MTT assay method.

#### Reference

- [1] Noordia, A., & Nurita, T. 2018 The Training of Aloe vera in the south of mulyorejo village. *ABDI Journal* **3** 2 84–87
- [2] Sujatha, G., Senthil Kumar, G., Muruganandan, J., & Srinivasa Prasad, T. 2014 Aloe vera in dentistry. *Journal of Clinical and Diagnostic Research* **8** 10 101-102
- [3] Kresnoadi, U. 2012 The increasing of fibroblast growth factor 2, osteocalcin, and osteoblast due to the induction of the combination of Aloe vera and 2% xenograft concelous bovine. *Dental Journal* **45** 4 228
- [4] Chantarawatit, P., Sangvanich, P., Banlunara, W., Soontornvipart, K., & Thunyakitpisal, P. 2014 Acemannan sponges stimulate alveolar bone, cementum and periodontal ligament regeneration in a canine class II furcation defect model. *Journal of Periodontal Research*, **49** 2 164–178
- [5] Beya, W., Davidson, B., & Erlwanger, K. H. 2012 The effects of crude aqueous and alcohol extracts of Aloe vera on growth and abdominal viscera of suckling rats. *African Journal of Traditional, Complementary and Alternative Medicines*, **9** 4 553–560.
- [6] Hu, Y., Xu, J., & Hu, Q. 2003 Evaluation of Antioxidant Potential of Aloe vera (*Aloe barbadensis* Miller) Extracts Evaluation of Antioxidant Potential of Aloe vera (*Aloe barbadensis* Miller) Extracts, *Journal Agric Food Chem*, **51** 7788–7791
- [7] Pangestuti, R., & Kim, S.-K. 2010 Neuroprotective Properties of Chitosan and Its Derivatives. *Marine Drugs*, **8** 7 2117–2128
- [8] Silva, S. S., Popa, E. G., Gomes, M. E., Cerqueira, M., Marques, A. P., Caridade, S. G., Reis, R. L. 2013 An investigation of the potential application of chitosan/aloe-based membranes for regenerative medicine. *Acta Biomaterialia*, **9** 6 6790–6797.
- [9] Ariani, M. D., Yuliati, A., & Adiarto, T. 2009 Toxicity testing of chitosan from tiger prawn shell waste on cell culture. *Dental Journal*, **42** 1 15–20.
- [10] Meizarini, A., Munadzirroh, E., & Rachmadi, P. 2015 The cytotoxicity of restorative material Cyanoacrylic with variation of powder and liquid ratio using MTT Assay, *Research Journal of Medika Eksakta*, **6** 1 16–25.
- [11] Fazwishni S, Hadijono BS. 2000 Cytotoxicity test using MTT Assay. *JKGUI* **7** 28–32
- [12] Yousef, H., Sharbeh, A., Ponnuraj, T., & Raja, R. A. 2012 In vitro cytotoxicity evaluation porcelain using human cell lines of dental, *The Journal of the School of Dental Science Universitas Sains Malaysia*, **7** 68–74.
- [13] Mestieri, L. B., Gomes-Cornélio, A. L., Rodrigues, E. M., Faria, G., Guerreiro-Tanomaru, J. M., & Tanomaru-Filho, M. 2017 Cytotoxicity and bioactivity of calcium silicate cements combined with niobium oxide in different cell lines. *Brazilian Dental Journal*, **28** 1 65–71.
- [14] Telli C, Serper A, Dogan AL, Guc D. 1999 Evaluation of the cytotoxicity of calcium phosphate root canal sealers by MTT Assay. *J Endodon*, **25** 8 3-11
- [15] Zeng Liantao, 2008 Absorption and distribution of chitosan in mice after oral administration. *J Carbohydrat Polymer* **7** 1 435-44
- [16] Rochima E, Suhartono MT, Syah D, Sugiono, 2007 The viscosity and molecular weight chitosan from chitin deacetylase enzymatic reaction Isolat *Bacillus Papandayan*. *Journal of Agricultural* 2-10
- [17] Silva, S. S., Caridade, S. G., Mano, J. F., & Reis, R. L. 2013 Effect of crosslinking in chitosan/aloe vera-based membranes for biomedical applications. *Carbohydrate Polymers*, **98** 1 581–588.
- [18] Srividya, S., Sastry, T. P., Jeevitha, D., & Samiksha, N. 2014 Synthesis and Characterization of a Novel Bone Graft Material. *International Journal of Drug Regulatory Affairs* 85–90.